

Institution: University of Maryland || Sign In as Member/Non-Member ||  
Contact Subscription Administrator at your institution || [FAQ](#)

Vol. 275, Issue 1, C1-C24, July 1998

## INVITED REVIEW

# Interaction of reactive oxygen species with ion transport mechanisms

Joseph I. Kourie

Membrane Transport Group, Department of Chemistry, The  
Faculties, The Australian National University, Canberra,  
Australian Capital Territory 0200, Australia

- ▶ [Abstract of this Article](#)
- ▶ [Reprint \(PDF\) Version of this Article](#)
- ▶ Similar articles found in:  
[Am J Physiol Cell Physiol Online](#)  
[PubMed](#)
- ▶ [PubMed Citation](#)
- ▶ This Article has been cited by:  
[other online articles](#)
- ▶ Search Medline for articles by:  
[Kourie, J. I.](#)
- ▶ Alert me when:  
[new articles cite this article](#)
- ▶ [Download to Citation Manager](#)

## ▶ ABSTRACT

The use of electrophysiological and molecular biology techniques has shed light on reactive oxygen species (ROS)-induced impairment of surface and internal membranes that control cellular signaling. These deleterious effects of ROS are due to their interaction with various ion transport proteins underlying the transmembrane signal transduction, namely, 1) ion channels, such as  $\text{Ca}^{2+}$  channels (including voltage-sensitive L-type  $\text{Ca}^{2+}$  currents, dihydropyridine receptor voltage sensors, ryanodine receptor  $\text{Ca}^{2+}$ -release channels, and D-myo-inositol 1,4,5-trisphosphate receptor  $\text{Ca}^{2+}$ -release channels),  $\text{K}^{+}$  channels (such as  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels, inward and outward  $\text{K}^{+}$  currents, and ATP-sensitive  $\text{K}^{+}$  channels),  $\text{Na}^{+}$  channels, and  $\text{Cl}^{-}$  channels; 2) ion pumps, such as sarcoplasmic reticulum and sarcolemmal  $\text{Ca}^{2+}$  pumps,  $\text{Na}^{+}$ - $\text{K}^{+}$ -ATPase ( $\text{Na}^{+}$  pump), and  $\text{H}^{+}$ -ATPase ( $\text{H}^{+}$  pump); 3) ion exchangers such as the  $\text{Na}^{+}/\text{Ca}^{2+}$  exchanger and  $\text{Na}^{+}/\text{H}^{+}$  exchanger; and 4) ion cotransporters such as  $\text{K}^{+}$ - $\text{Cl}^{-}$ ,  $\text{Na}^{+}$ - $\text{K}^{+}$ - $\text{Cl}^{-}$ , and  $\text{P}_i$ - $\text{Na}^{+}$  cotransporters. The mechanism of ROS-induced modifications in ion transport pathways involves 1) oxidation of sulfhydryl groups located on the ion transport proteins, 2) peroxidation of membrane phospholipids, and 3) inhibition of membrane-bound regulatory enzymes and modification of the oxidative phosphorylation and ATP levels. Alterations in the ion transport mechanisms lead to changes in a second messenger system, primarily  $\text{Ca}^{2+}$  homeostasis, which further augment the abnormal electrical activity and distortion of signal transduction, causing cell dysfunction, which underlies pathological conditions.


ischemia-reperfusion; muscle pathologies; thiol group; calcium homeostasis; membrane compartmentation; reducing and oxidizing agents

## ▶ INTRODUCTION

- ▲ [Top](#)
- [Abstract](#)
- ▼ [Introduction](#)
- ▼ [Conclusions](#)
- ▼ [References](#)




(Fenton reaction)



Experimentally, different ROS-generating and/or ROS-identifying systems have been used to examine the ROS-induced modifications of ion transport pathways (see Tables 1-9). These include 1)  $\text{H}_2\text{O}_2$ ; 2) *tert*-butyl hydroperoxide (*t*-BHP), a substrate of glutathione peroxidase; 3) *t*-butoxy ( $\text{RO}\cdot$ ) or *t*-butylperoxy ( $\text{ROO}\cdot$ ) radical-generating systems; 4) hypoxanthine (HX)/XO, a source for  $\text{O}_2^{\cdot -}$ ,  $\text{H}_2\text{O}_2$ , and  $\cdot\text{OH}$  production; 5) dihydroxyfumaric acid (DHF); 6) cumene hydroperoxide or purine/XO; 7) photooxidizing rose bengal, a source for  $^1\text{O}_2$ ; 8) ionizing  $\gamma$ -irradiation, diethylenetriaminepentaacetic acid, and catalase/XO; 9) HX/XO,  $\text{FeCl}_3$ , and ADP; 10) phorbol myristate acetate activation of  $\text{H}_2\text{O}_2$  production in neutrophils; and 11) the free radical scavenger *N*-acetyl-L-cysteine. Pharmacological identification and dissection of the combined ROS effects are achieved by examining the modulatory effects of specific scavengers for  $\text{H}_2\text{O}_2$ ,  $\text{O}_2^{\cdot -}$ ,  $\cdot\text{OH}$ , and  $^1\text{O}_2$  such as catalase, superoxide dismutase (SOD), desferrioxamine, and histidine, respectively. It is thought that the effects of the non-free-radical  $\text{H}_2\text{O}_2$  are caused by producing more highly reactive oxygen species such as the free radicals  $\text{O}_2^{\cdot -}$  and  $\cdot\text{OH}$ . In particular,  $\cdot\text{OH}$  reacts rapidly with many substances, e.g., DNA, lipid, and carbohydrates.

A flowchart of the major processes of ROS underlying pathologies is shown in Fig. 1. The pathologies that have been attributed to ROS-induced cell dysfunction include 1) cardiac stunning and arrhythmia (see Refs. 35 and 61); 2) skeletal muscle injury (see Refs. 130 and 151); 3) neurological conditions (see Refs. 91 and 126), e.g., neuronal damage in Parkinson's disease (see Ref. 27); 4) neurotoxicity (107); 5) Alzheimer's disease (see Refs. 6 and 171); 6) diabetes (see Ref. 123), apoptosis of T lymphocytes (see Ref. 37), and gastric mucosal injury (see Ref. 160); and 7) hypertension (156). Some of these effects can be suppressed by free radical scavengers (9, 54, 94, 104).



**Fig. 1.** Flow chart of major processes of reactive oxygen species (ROS) pathologies.

[View larger version \(29K\):](#)

[\[in this window\]](#)

[\[in a new window\]](#)

In skeletal muscle, exercise increases the rate of ROS production (30, 130, 158). The enhancement of ROS production due to the increase in activity of mitochondrial electron carriers, low catalase concentrations, the sudden changes in oxygen supply and consumption, and the presence of high levels of myoglobin acting as a catalyst for the formation of oxidants is thought to cause skeletal muscle injury (see Ref. 130). The increase of free radicals in skeletal muscle and liver cells during exhaustive exercise is associated with a decrease in mitochondrial respiratory control, loss of sarcoplasmic reticulum (SR)/endoplasmic reticulum (ER) integrity, and increased levels of peroxidation products and lipid peroxidation. These effects are similar to those observed in vitamin E-deficient animals (30).

## ► ROS INTERACTION WITH ION TRANSPORT PATHWAYS


The interaction of ROS with ion transport pathways in muscles can be deduced indirectly from changes in their membrane properties. Cosentino et al. (29) demonstrated the role of  $O_2^-$  in the mediation of endothelium-dependent contraction. It has also been demonstrated that  $H_2O_2$  potentiates twitch tension in cardiac (84, 136) and skeletal muscles (124, 136), and this induced tension can be decreased by catalase, a specific enzyme that hydrates  $H_2O_2$  (136). The effects are usually characterized by amplification of tension and tension oscillation, followed by spontaneous contractions (84, 124). This effect of  $H_2O_2$  is not mediated via end effects on the myofilaments (111, 124). This suggests that the signal transduction pathways are affected by ROS. Early studies revealed that the effects of ROS on membrane properties could be deduced from electrophysiological parameters of the membrane. These include changes in membrane current and potential, ionic gradients, action potential duration and amplitude, afterdepolarization, and spontaneous activity and loss of excitability (see Refs. 40, 166, 167).

The effects of ROS-generating systems on membrane potential are now well established. It has been

demonstrated that X/XO as a ROS-generating system caused membrane depolarization and a decrease in the action potential amplitude and maximum rate of rise of action potentials in guinea pig ventricular myocardium (127). Delayed afterdepolarization and early afterdepolarization induced by *t*-BHP, DHF, and X/XO in guinea pig papillary muscle and canine ventricular myocytes have also been demonstrated (4, 5, 122). ROS-induced membrane depolarization has been attributed to inhibition of a  $\text{Na}^+$  current (11) or an inward  $\text{K}^+$  current (121), activation of an inwardly directed nonselective cation current (115, 152), and increase in a  $\text{Ca}^{2+}$  current that is associated with changes in intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ). Similarly, the oscillation in  $[\text{Ca}^{2+}]_i$  has been implicated in arrhythmic afterdepolarization (113).

ROS-induced shortening of the action potential duration has been attributed to a possible increase in a delayed rectifying  $\text{K}^+$  current and decrease in activation of ATP-sensitive  $\text{K}^+$  ( $\text{K}_{\text{ATP}}$ ) channels and  $\text{Ca}^{2+}$  currents (121, 146). Exogenous ROS-induced changes in the electromechanical function and metabolism in isolated rabbit and guinea pig ventricles shortened the duration of the action potential, indicating a decrease in the  $\text{Ca}^{2+}$  current and time-dependent outward current (50). More recently, Tokube et al. (169) reported biphasic changes in the action potential duration, with initial lengthening of the action potential due to a rapid decrease in whole cell  $\text{K}^+$  currents and subsequent shortening due to a decrease of whole cell  $\text{Ca}^{2+}$  current and increase in the single ATP-sensitive time-dependent outward  $\text{K}^+$  current.

In cardiac, smooth, and skeletal muscles the deleterious effects of ROS, produced by leaked electrons from the electron transport system of the mitochondria, are due to their interaction with various ion transport proteins underlying transmembrane signal transduction (Fig. 2). Figure 2 indicates that an important feature of ROS interaction with ion transport proteins is the modification in  $\text{Ca}^{2+}$  homeostasis that ultimately causes muscle pathologies.



View larger version (22K):  
[\[in this window\]](#)  
[\[in a new window\]](#)

**Fig. 2.** Generalized scheme showing ROS-modulated ion transport mechanisms that control  $\text{Ca}^{2+}$  homeostasis ( $\text{Ca}^{2+}$  pool) in muscles. These transport mechanisms include 1) ion channels:  $\text{Ca}^{2+}$  channels, including voltage-sensitive L-type  $\text{Ca}^{2+}$  currents ( $\text{V Ca}^{2+}$ ), ligand  $\text{Ca}^{2+}$  channels ( $\text{R Ca}^{2+}$ ), dihydropyridine receptor (DHPR) voltage sensor, ryanodine receptor (RyR)  $\text{Ca}^{2+}$ -release channels, and D-myo-inositol 1,4,5-trisphosphate receptor ( $\text{IP}_3\text{-R}$ )  $\text{Ca}^{2+}$ -release channels;  $\text{K}^+$  channels, such as ATP-sensitive  $\text{K}^+$  channels; and  $\text{Cl}^-$  channels, such as the small  $\text{Cl}^-$  (SCL) channel; 2) ion pumps: such as sarcoplasmic reticulum (SR) and sarcolemmal (S) ATP  $\text{Ca}^{2+}$  pumps and  $\text{Na}^+\text{-K}^+\text{-ATPase}$  ( $\text{Na}^+$  pump); and 3) ion exchangers:  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. Excited cell membrane (sarcolemma of skeletal muscle) or specific receptor (R, in sarcolemma of cardiac and smooth muscles) communicate with the  $\text{Ca}^{2+}$  sender (SR in skeletal and smooth muscles and sarcolemma in cardiac muscle) by means of the T tubule (T-T) or by second messengers, such as cAMP (cardiac muscle) or  $\text{IP}_3$  (smooth muscle). M, mitochondria; MF, myofilaments; R, receptor  $\beta$  (cardiac muscle) and  $\alpha_1$  (smooth muscles).

## Ion Channels

**$\text{Ca}^{2+}$  channels.** L-TYPE VOLTAGE-SENSITIVE  $\text{Ca}^{2+}$  CURRENTS. L-type voltage-sensitive  $\text{Ca}^{2+}$  channels play an important role in  $\text{Ca}^{2+}$  homeostasis in ventricular myocytes. Hence numerous studies have been conducted to examine the effects of ROS on these channels and to determine their contribution to the alterations in  $\text{Ca}^{2+}$  homeostasis under adverse conditions (Table 1). It appears that the data for the effects of ROS on current peak, amount of current, and kinetics of L-type  $\text{Ca}^{2+}$  channels in ventricular myocytes are conflicting. It has also been reported that  $\text{H}_2\text{O}_2$  has no influence on L-type  $\text{Ca}^{2+}$  current (100, 101).

In contrast to the finding that ROS-induced reduction in peak current was associated with an increase in mean current due to slowing of the inactivation (26), Tokube et al. (169) reported a decrease in the current peak with no changes in the activation time course of this current. On the other hand, Cerbai et al. (20), Matsuura and Shattock (115), and Moghadam and Winlow (119) reported a decrease in L-type  $\text{Ca}^{2+}$  current. This decrease has been attributed to  $\text{Ca}^{2+}$ -induced channel inactivation (see Ref. 115). The  $\text{H}_2\text{O}_2$ -induced decrease in the inward  $\text{Ca}^{2+}$  current in cultured *Lymanaea* neurons is dose dependent (119). There are data suggesting that overload due to  $\text{Ca}^{2+}$  influx through the voltage-gated  $\text{Ca}^{2+}$  channel can be ruled out, since free radicals and  $\text{H}_2\text{O}_2$  inhibit the voltage-sensitive L-type  $\text{Ca}^{2+}$  current (48, 50, 51, 121). The inhibitory effects of HX/XO and DHF as ROS-generating systems were reversed with SOD and catalase, suggesting that both  $\text{O}_2^-$  and  $\text{H}_2\text{O}_2$  are effective (Table 1), whereas the effects of the

cumene/XO ROS-generating system were irreversible (48). Internal oxidative agents used on ion channels also show that 4,4'-dithiodipyridine [DTDP; a lipophilic sulfhydryl (SH)-oxidizing agent] and thimerosal {[*o*-carboxyphenyl]thio}ethyl mercury sodium salt, a hydrophilic SH-oxidizing agent} inhibit the activity of cloned rabbit smooth muscle L-type  $\text{Ca}^{2+}$  channels (23).

**Table 1. Effects of ROS on DHPR and L-type  $\text{Ca}^{2+}$  currents**

**View this table:**

[\[in this window\]](#)

[\[in a new window\]](#)

$\text{Ca}^{2+}$  channel blockers have been used to identify the  $\text{Ca}^{2+}$  pathway contributing to changes in cytoplasmic  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_{\text{cyt}}$ ). The  $\text{Ca}^{2+}$  channel blocker nifedipine blocks  $\text{O}_2^-$ -induced increases in  $[\text{Ca}^{2+}]_{\text{cyt}}$  in human myometrial cells (112). Although indirect and inconclusive, the finding is taken by the authors to indicate that the increase in  $\text{Ca}^{2+}$  is mediated via an  $\text{O}_2^-$ -affected voltage-sensitive L-type  $\text{Ca}^{2+}$  channel. Recently, Ueda et al. (171) reported that free radicals, monitored with 2',7'-dichlorofluorescein diacetate, may be involved in amyloid protein potentiation of  $\text{Ca}^{2+}$  influx through L-type voltage-sensitive  $\text{Ca}^{2+}$  channels. Amyloid  $\beta$  protein, which accumulates in the brain of Alzheimer patients, generates  $\text{Ca}^{2+}$ -independent free radicals that potentiate the influx of  $\text{Ca}^{2+}$  through L-type voltage-sensitive  $\text{Ca}^{2+}$  channels in rat cultured cortical and hippocampal neurons. The neurotoxicity (see Refs. 6 and 170) caused by this influx is attenuated by nimodipine (171, 180) and vitamin E (171).

**DIHYDROPYRIDINE RECEPTOR VOLTAGE SENSOR.** There is indirect evidence for the effect of ROS on the dihydropyridine receptor (DHPR). It has been found that  $\text{H}_2\text{O}_2$  prevents  $\text{Ag}^+$  contractions and  $\text{Ag}^+$  inhibition of excitation-contraction (E-C) coupling in single skeletal muscle fibers from *Rana temporaria* or *R. catesbeiana* (124). Recently, Oba et al. (124) proposed that  $\text{H}_2\text{O}_2$  induces skeletal muscle dysfunction by acting on the DHPR and the ryanodine receptor (RyR) in T tubule and SR, respectively. Tension experiments on skinned single muscle fibers from *R. catesbeiana* reveal that 1.5-6 mM  $\text{H}_2\text{O}_2$  potentiates decaying twitches indicative of a direct action on the DHPR, although neither resting nor action potentials were affected (124). Decaying twitches were seen in the presence of 5 mM dithiothreitol (DTT) and were amplified and slowed with BAY K 8644. Binding studies that also indicate a ROS-induced decrease in this current may suggest a direct effect on the channel protein. Kaneko et al. (87) observed a reduction in DHP binding sites in the membranes of heart cells exposed to oxygen free radicals. Similarly, in guinea pig ventricular myocytes, the ROS-generating system DHF reduced  $[\text{^3H}]\text{PN-200-110}$  binding sites of DHP, underlay the observed reduction in L-type  $\text{Ca}^{2+}$  currents, and was prevented by SOD and catalase (58). These authors postulated that these changes in the DHPRs, which reduce  $\text{Ca}^{2+}$  currents, mediate the mechanical dysfunction associated with oxidative stress. However, the effects of  $\text{H}_2\text{O}_2$  or other ROS on single DHPR channel activity have not been reported yet.

**RYR  $\text{Ca}^{2+}$ -RELEASE CHANNELS.** In cardiac and skeletal muscles the RyR  $\text{Ca}^{2+}$ -release channels are essential in maintaining  $\text{Ca}^{2+}$  homeostasis that underlies the mechanism of muscle contraction and relaxation. There are only a few studies regarding the effects of ROS on these channels. However, these studies have established that the effect of ROS on  $\text{Ca}^{2+}$  homeostasis can be attributed, in part, to  $\text{Ca}^{2+}$  release from the SR. There is also biochemical evidence revealing that ROS modify the structure and function of the cardiac SR RyR  $\text{Ca}^{2+}$ -release channel, where the initial increase in the probability of the channel being in the open state ( $P_o$ ) is followed by irreversible loss of the channel function (69). In skeletal muscle,  $\text{H}_2\text{O}_2$  induces SR  $\text{Ca}^{2+}$  release that can be enhanced with  $\text{Cu}^{2+}$  (170), probably through skeletal SR RyR  $\text{Ca}^{2+}$ -release channels (124). Similarly, in sheep cardiac SR,  $\text{H}_2\text{O}_2$  (3-5 mM) directly modified the gating of the RyR  $\text{Ca}^{2+}$ -release channel, causing an increase in the  $P_o$ , without affecting the conductance or channel modulation with ATP, caffeine,  $\text{Mg}^{2+}$ , or ryanodine (12). It appears that  $\text{Ca}^{2+}$  release from the SR can be induced using different ROS-generating systems (Table 2). For example, a 106-kDa  $\text{Ca}^{2+}$ -release channel protein from the SR of skeletal muscle is also activated by rose bengal (185). More recently, using the  $\text{Ca}^{2+}$  sensitivity and the maximum  $\text{Ca}^{2+}$ -activated force of isolated skinned fibers as indirect investigative parameters, Posterino and Lamb (132) reported that reducing agents do not inhibit the E-C coupling and that oxidizing agents do not cause a significant  $\text{Ca}^{2+}$  release under physiological conditions. However, in the absence of supporting data at the single  $\text{Ca}^{2+}$  channel and ion pump levels, it is difficult to determine any direct modifications in the  $\text{Ca}^{2+}$ -transport pathways or other mechanisms of  $\text{Ca}^{2+}$  release and uptake.

**Table 2. Effects of ROS on RyR  $\text{Ca}^{2+}$ -release channels**

**View this table:**  
[\[in this window\]](#)  
[\[in a new window\]](#)

**D-MYO-INOSITOL 1,4,5-TRISPHOSPHATE RECEPTOR  $\text{Ca}^{2+}$ -RELEASE CHANNELS.** ROS-induced  $\text{Ca}^{2+}$  release via modifications in D-myoinositol 1,4,5-trisphosphate ( $\text{IP}_3$ )-induced  $\text{Ca}^{2+}$  release at the single-channel level remains to be experimentally observed. However, it has been reported that  $\text{O}_2^-$  stimulates  $\text{IP}_3$ -induced  $\text{Ca}^{2+}$  release from the SR of vascular smooth muscle (165). Furthermore, it has been proposed that SH reagents may induce  $\text{Ca}^{2+}$  release by sensitizing the  $\text{IP}_3$   $\text{Ca}^{2+}$ -release receptor (118). The data reported by Elmoselhi et al. (43) indicate that oxygen free radicals modify  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  channels (see *Changes in  $\text{Ca}^{2+}$  Homeostasis*). It has also been reported that oxidized glutathione (GSSH) decreases the luminal  $\text{Ca}^{2+}$  content of the endothelial cell line  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  store (64).

**$\text{K}^+$  channels.  $\text{Ca}^{2+}$ -ACTIVATED  $\text{K}^+$  CHANNELS.** The role of ROS in modulating ion channels has also been inferred from the use of ion channel blockers together with ROS-generating systems. The  $\text{K}^+$  channel blocker quinidine hydrochloride reduced  $\text{Ca}^{2+}$ -dependent chemiluminescence products, indicative of oxygen radical production, in human eosinophils (143). They postulated that production of oxygen free



radicals by the membrane-bound NADPH oxidase may be mediated by  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  ( $\text{K}_{\text{Ca}}$ ) channels in human eosinophils and that this mechanism may underlie the role of eosinophils in the pathogenesis of allergic diseases. Relaxation evoked by nonneurogenic electrical field stimulation, via generation of free radicals, also modified  $\text{Ca}^{2+}$ -dependent channels (1, 81, 186). In contrast to the  $\text{H}_2\text{O}_2$ -induced reversible inhibition [with DTT and reduced glutathione (GSH)] of  $\text{K}_{\text{Ca}}$  channels in the plasma membrane of bovine aortic endothelial cells (18), the large  $\text{K}_{\text{Ca}}$  channel in skeletal muscle from mouse is insensitive to as high as 50 mM  $\text{H}_2\text{O}_2$  concentration (179). Differences in  $\text{H}_2\text{O}_2$ -induced modification in channel activity could be attributed to difference in tissue types. For example, it has been found that reducing agents decrease the activity of  $\text{K}_{\text{Ca}}$  channels in pulmonary, but not in ear, arterial smooth muscle cells of rabbit (128). The significance of  $\text{H}_2\text{O}_2$  inhibition of  $\text{K}_{\text{Ca}}$  channels derives from the fact that disruption of  $\text{Ca}^{2+}$  homeostasis is mediated via depolarization of the membrane potential (see Table 3) (18).

**Table 3.** Effects of ROS on  $\text{K}_{\text{Ca}}$  channels

**View this table:**

[\[in this window\]](#)

[\[in a new window\]](#)

INWARD AND OUTWARD  $\text{K}^+$  CURRENTS. The modifications in ion channels that underlie ROS-induced changes in the duration of action potential in cardiac cells include, in addition to the ROS-induced decrease in L-type  $\text{Ca}^{2+}$  current, a suppression of the delayed outward and the inward  $\text{K}^+$  currents (77, 169). Electrically evoked relaxation, which generates free radicals in canine airway smooth muscle relaxation, was not sensitive to removal of external  $\text{K}^+$  but was sensitive to tetraethylammonium, high KCl concentrations, charybdotoxin, quinine, and free radicals (186). This was deduced from the action of the free radical scavenger *N*-acetylcysteine and was mimicked by  $\text{H}_2\text{O}_2$ , whereas SOD and catalase were ineffective. In guinea pig ventricular myocytes the ROS-generating system DHF also reduced the outward  $\text{K}^+$  current that determines the prolongation of the action potential as a result of exposure to oxygen free radicals (20, 58). Similarly, Kuo et al. (104) found that nonlethal ionizing  $\gamma$ -irradiation (10 cGy) transiently ( $t_{1/2}$  90 min) induced whole cell voltage-dependent outward  $\text{K}^+$  currents, mimicking  $\text{H}_2\text{O}_2$  and heat stress in activating this current, with no changes in membrane potential of  $-70$  mV, intracellular  $\text{K}^+$  concentration ( $[\text{K}^+]_i$ ), or ATP levels.

Several inward and outward  $\text{K}^+$  channels are affected by different ROS-generating systems (Table 4). In guinea pig cardiac ventricular myocytes HX/XO (ROS production that is indicated from adrenochrome formation from adrenaline) decreased the inward  $\text{K}^+$  current (26). In rabbit sinoatrial atrioventricular node preparation, *t*-BHP transiently increased the spontaneous firing frequency, increased the amplitude of the action potential, and induced biphasic changes in  $\text{Ca}^{2+}$  current, delayed rectifying  $\text{K}^+$  current, and hyperpolarization-activated inward current (145). In guinea pig ventricular cells cumene hydroperoxide decreased whole cell inward rectifier  $\text{K}^+$  current and inhibited the single inward rectifier  $\text{K}^+$  channel

without altering the unit amplitude of single-channel current (121). In *Xenopus* oocytes  $H_2O_2$  reversibly and specifically inhibited the time-dependent fast activation of a certain voltage-gated  $K^+$  channel concomitantly associated with an increase in  $K^+$  currents of cloned  $K^+$  channels KShIIC, KShIID, and HukII. Other cloned voltage-dependent channels were not affected by 1.6 mM  $H_2O_2$ , e.g., *Shaker* 29-4 and KShIIIA.1 (173). Recently, Dupart et al. (37) found that photoactivation of rose bengal induced inhibition of the cloned  $K^+$  channel activity of *Shaker* channels Kv1.3, Kv1.4, and Kv1.5, *Shaw* channel Kv3.4, and inward  $K^+$  rectifier IRK3, whereas *Shaker* Kv1.2, *Shab* channels Kv2.1 and Kv2.2, *Shal* channel Kv4.1, and inward rectifiers IRK1, ROMK1, and hIsK were not affected. On the other hand, *t*-BHP removed the fast inactivation of the *Shaker*  $K^+$  channels Kv1.4 and Kv3.4, whereas other  $K^+$  channels were not affected. These findings indicate that ROS effects depend on the ROS-generating system as well as on the type of channel protein under investigation.

**Table 4.** Effects of ROS on inward, outward and other  $K^+$  currents

**View this table:**

[\[in this window\]](#)

[\[in a new window\]](#)

ATP-SENSITIVE  $K^+$  CHANNELS. The effects of  $H_2O_2$  and HX/XO on  $K_{ATP}$  channels in cardiac and pancreatic cells have been reported (Table 5). Their effects on this channel are reminiscent of the effects of hypoxic conditions. For example, hypoxic conditions induced a time-independent  $K^+$  current through  $K_{ATP}$  in isolated heart cells of the guinea pig (7). Similarly, in guinea pig ventricular myocyte cells, the HX/XO ROS-generating system increased the  $P_o$  of  $K_{ATP}$  and glibenclamide-sensitive  $K^+$  channels (77, 169). Concentrations of  $>30 \mu M H_2O_2$  also increased the activity of  $K_{ATP}$  channels in the plasma membrane of rat pancreatic  $\beta$ -cells in whole cell current of perforated cells but not that of conventional whole cell configuration. Furthermore, it increased single-channel activity in the cell-attached configuration but not in the inside-out configuration. This effect was inhibited with tolbutamide, glyceraldehyde, and 2-ketoisocaproic acid (123). Evidence for direct effects of  $H_2O_2$  on  $K_{ATP}$  channels can be deduced from studies where ROS effects were examined on excised membrane patches. For example, Ichinari et al. (73) observed a dose-dependent  $H_2O_2$ -induced increase in  $P_o$  of the  $K_{ATP}$  channel in the isolated inside-out configuration. It has been proposed that a  $H_2O_2$ -induced increase in  $K_{ATP}$  channel activity in  $\beta$  cells that remained sensitive to ATP was due to indirect channel opening via inhibition of glycolysis and/or oxidative phosphorylation, leading to a decrease in the cytosolic concentration of ATP (123), as previously suggested for the  $H_2O_2$ -induced increase in  $K_{ATP}$  channels in guinea pig ventricular myocytes (50). In  $\beta$  cells,  $H_2O_2$  stimulated  $K_{ATP}$  currents but not  $Ca^{2+}$  current (100, 101). They proposed that the SH-oxidizing agents 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) and  $H_2O_2$  may act 1) on different SH targets or 2) via a mechanism other than oxidation of SH groups.  $H_2O_2$ -induced relaxation in rabbit airway smooth muscle has been attributed to the activation of

an  $\alpha$ -catalytic subunit with a relative molecular weight ( $M_r$ ) of 90,000-110,000 and a  $\beta$ -subunit with an  $M_r$  of 40,000-60,000. The  $\text{Na}^+$  pump is important for maintaining coronary tone. The pump transports three  $\text{Na}^+$  and two  $\text{K}^+$  per one ATP hydrolyzed against their concentration gradients, generating internal negative charges. The intracellular/extracellular concentrations in muscle cells are typically 5-115/130 mM  $\text{Na}^+$  and 130/5 mM  $\text{K}^+$  (see Ref. 42). There are several reaction steps in the function of the pump. The hydrolytic and transport reactions of the pump can be uncoupled. For example, treatment of inside-out erythrocyte vesicles with trypsin or chymotrypsin uncouples the transport of  $\text{Na}^+$  from the  $\text{Na}^+$ - $\text{K}^+$ -ATPase (62).  $\text{K}^+$ -activated ouabain-sensitive *p*-nitrophenylphosphatase is associated with the hydrolytic step of the  $\text{Na}^+$  pump. Conflicting effects of ROS on  $\text{Na}^+$  pumps have been reported (see Ref. 62). Some of these differences in ROS effects are due to the methods used for ROS generation, in which different individual ROS may not act via the same mechanism (Table 8). These differences could also arise from resolution limitations of the flux experiments, which can be overwhelmed by the back flux of ROS-enhanced activated  $\text{K}^+$  channels.

**Table 8.** Effects of ROS on  $\text{Na}^+$ - $\text{K}^+$ -ATPase ( $\text{Na}^+$  pump)

**View this table:**

[\[in this window\]](#)

[\[in a new window\]](#)

Kim and Akera (90) examined  $\text{Na}^+$ - $\text{K}^+$ -ATPase or the  $\text{Na}^+$  pump in the sarcolemma of guinea pigs.  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity was deduced from ouabain-sensitive ATPase and was estimated from ouabain-sensitive  $^{86}\text{Rb}^+$  uptake. They found that scavengers (100 U/ml SOD, 150 U/ml catalase, 50 mM DMSO, 10 mM histidine, and 50  $\mu\text{g}/\text{ml}$  vitamin E or 1 mM XO inhibitor allopurinol) of  $\text{O}_2$  free radicals inhibited ischemia and reperfusion-induced reduction in  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity and specific [ $^3\text{H}$ ]ouabain binding. The pump also appeared to be sensitive to changes in membrane phospholipids. It has been shown that  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity is also inhibited by lipoxygenase, dihydroxyfumarate, ascorbate- $\text{FeCl}_2$ , and cumene hydroperoxide (15, 99, 163). Jamme et al. (80) reported that the inhibition of mouse cerebral  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity by ultraviolet C (UV-C)-generated  $\cdot\text{OH}$  and peroxy ( $\text{ROO}\cdot$ ) radicals is mediated via lipid peroxidation-induced disruption of membrane integrity that consequently results in conformational changes, leading to inactivation of membrane-bound proteins. It has also been suggested that ascorbate- $\text{FeCl}_2$ -induced inactivation of cerebral  $\text{Na}^+$ - $\text{K}^+$ -ATPase is due to lipid peroxidation-induced reduction in the affinity for  $\text{Na}^+$  and  $\text{K}^+$  and an increase in ATP and ouabain affinities (117). The inhibitory effects of iron-generated free radicals on the activity of  $\text{Na}^+$ - $\text{K}^+$ -ATPase can be reversed by antioxidants (138).

It appears that the effects of free oxide radicals during ischemia and/or perfusion on  $\text{Na}^+$ - $\text{K}^+$ -ATPase are not due to depletion of ATP, since ATP recovers on reperfusion while the free oxide radicals continue to enhance the decrease in  $\text{Na}^+$ - $\text{K}^+$ -ATPase and glycoside binding sites (see Refs. 82 and 90). The effects of X/XO on cardiac  $\text{Na}^+$ - $\text{K}^+$ -ATPase are not clear. Vlessis et al. (175) reported inhibition of both  $\text{Na}^+$

transport and the  $\text{Na}^+\text{-K}^+\text{-ATPase}$ , whereas Kukreja et al. (103) found that  $\text{X/XO}$  was ineffective while  $\text{Na}^+\text{-K}^+\text{-ATPase}$  was inactivated by  $\text{H}_2\text{O}_2$ ,  $\text{HClO}_4$ , and  $\text{NH}_2\text{Cl}$  treatments. Vinnikova et al. (174) found that the  $^1\text{O}_2$ -induced inhibitory effect on  $\text{Na}^+\text{-K}^+\text{-ATPase}$  was prevented by the  $^1\text{O}_2$  scavenger, histidine, whereas SOD, catalase, and mannitol were not effective in providing such protection. Therefore, these data rule out inhibitory effects due to  $\text{O}_2^-$ ,  $\text{H}_2\text{O}_2$ , and  $\cdot\text{OH}$ . Elmoselhi et al. (42) suggested that  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$  uncouple the hydrolytic activity of the  $\text{Na}^+$  pump from  $\text{Rb}^+(\text{K}^+)$  uptake. They proposed that such uncoupling under ischemic conditions and reperfusion would damage coronary artery smooth muscle as a result of continuous ionic imbalance and starvation of the cell via continuous ATP hydrolysis. The levels of  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$  required for uncoupling the  $\text{Na}^+$  pump are higher than those affecting other processes; hence  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$  effects are unlikely to be directly due to uncoupling of the hydrolytic and transport reactions of the  $\text{Na}^+$  pump. It is not known whether ROS affect the various  $\text{Na}^+$  pump isoforms. For example, these isoforms differ in their  $\alpha$ - and  $\beta$ -subunits and in their affinities to  $\text{Na}^+$  and  $\text{K}^+$  (42). Differences in responses of these isoforms to specific ROS may shed light on their molecular mechanisms of action at the subunit level. In this regard it has been found that the  $\alpha_1$  and  $\alpha_2$  isoforms of the  $\text{Na}^+\text{-K}^+\text{-ATPase}$  differ in their sensitivities to oxidants (71, 184).

*$\text{H}^+$  pump.* The  $\text{H}^+$  pump is important for preventing a drastic intracellular acidification and for charge balance and membrane polarization. Oxidant stress-induced pH changes in peritoneal macrophages have been attributed to modifications in the plasmalemmal  $\text{H}^+\text{-ATPase}$  (see Ref. 14).

*Adenine nucleotide translocator, phosphate carrier, and uncoupling proteins.* These proteins are present in mitochondria. The phosphate carriers catalyze the electroneutral exchange of phosphate for hydroxyl ion. The adenine nucleotide carrier binds and transports adenine nucleotides, whereas uncoupling proteins bind purine nucleotides but transport  $\text{H}^+$ ,  $\text{OH}^-$ , or  $\text{Cl}^-$  (see Ref. 137). The effects of ROS on these protein transporters are not known. However, it is very likely that they are affected by ROS. First, it is known that in skeletal muscle and liver cells free radicals increase during exhaustive exercise and this increase is associated with a decrease in mitochondrial respiratory control (30). Second, ROS have deleterious effects on mitochondrial metabolism (see Ref. 50) and are linked to a leakage of electrons from mitochondria (see Ref. 91). Third, the presence of the SH groups on the cysteine residues and the SH-induced modification in permeation of phosphate,  $\text{Cl}^-$ , and  $\text{H}^+$  (137) also suggest that ROS may modify these transporters.

### *Ion Exchangers*

*$\text{Na}^+/\text{Ca}^{2+}$  exchanger.* The  $\text{Na}^+/\text{Ca}^{2+}$  exchanger couples the transport of three  $\text{Na}^+$  to that of a single  $\text{Ca}^{2+}$  in the opposite direction in two consecutive, yet separate steps (see Ref. 28). The  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, together with  $\text{Ca}^{2+}\text{-ATPase}$  of the ER/SR, regulates  $\text{Ca}^{2+}$  levels that underlie muscle contractility behavior under both normal and ischemic conditions (see Ref. 13). In cardiac muscle the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger contributes to force development, in particular, under glycosidic conditions (see Ref.



reabsorption of  $\text{NaCl}$  and  $\text{NaHCO}_3$ . There is little information on the effects of ROS on these exchangers in epithelial cells. ROS have been implicated in the increased activity of the cellular  $\text{Na}^+/\text{H}^+$  exchanger that is activated by phosphorylation in vascular myocytes from hypertensive rats (156). It has also been reported that exposure of human neutrophils to 100 nM *N*-formyl-methionyl-leucyl-phenylalanine activated the amiloride-sensitive  $\text{Na}^+/\text{H}^+$  exchanger, leading to an increase in intracellular pH from 7.22 to 7.8 (157). The ROS-generating system X/XO inhibited this transport system in isolated myocytes of rat heart and in sealed sarcolemmal vesicles of bovine heart (184), and the inhibition was reversed with catalase and SOD and, therefore, indicated that  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$  were the effective moieties. The effect of ROS on the  $\text{Cl}^-/\text{HCO}_3^-$  exchanger is unknown.

### *Ion Cotransporters*

**Cation- $\text{Cl}^-$  transporters.** The electroneutral transporters have important physiological roles, such as regulatory volume decrease and transepithelial salt transport (see Ref. 120). Their activity depends on the presence of all the transported ions. However, they differ pharmacologically with respect to the identity and stoichiometry of the transported ions. There is little direct or indirect information on the effect of ROS on these transporters.

**$\text{K}^+-\text{Cl}^-$  COTransport.** This cotransport system could also be modulated by ROS, since it has been reported that  $\text{K}^+-\text{Cl}^-$  cotransport in erythrocytes is modulated by SH groups. It is activated through *N*-ethylmaleimide (NEM)-induced SH alkylation and methylmethane thiosulfonate- or diamide-induced SH oxidation (see Ref. 187). It has also been found that phenazine methosulfate, a generator of oxygen free radicals, stimulated the reversible  $\text{K}^+-\text{Cl}^-$  cotransport system in human erythrocyte membranes (60).

**$\text{Na}^+-\text{K}^+-\text{Cl}^-$  COTransport.** The effects of oxidant stress, induced via cell incubation in *t*-BHP in the presence of bumetanide, show a decrease in the inward movement of  $\text{Rb}^+$ , indicating inhibition of the bumetanide-sensitive  $^{86}\text{Rb}^+$  pathway, which represented  $\text{Na}^+-\text{K}^+-\text{Cl}^-$  cotransport (41). Similarly, *t*-BHP inhibited  $\text{Na}^+-\text{K}^+-\text{Cl}^-$  cotransport in skeletal muscle (151).

**Other cotransporters.**  **$\text{Na}^+-\text{P}_i$  COTransport.**  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$  inhibited the  $\text{Na}^+-\text{P}_i$  transport system in isolated myocytes of rat heart and in sealed sarcolemmal vesicles of bovine heart (184). Furthermore, this ROS-induced inhibition was reversed with catalase and SOD. The effect of ROS on  $\text{Na}^+-\text{HCO}_3^-$  and  $\text{K}^+-\text{HCO}_3^-$  cotransporters is unknown.

## ► THE PRIMARY TRANSPORT PATHWAY AS A TARGET FOR ROS


ROS-induced changes in membrane properties are considered early events in response to oxidative stress. However, the molecular mechanism(s) for ROS action on ion transport pathways is not known. Hypothetically, the effects of ROS can be caused via direct effects on ion transport proteins. Ion channels

that have been thought to be a prime ROS target include a 106-kDa  $\text{Ca}^{2+}$ -release channel (162, 185), DHPR and RyR  $\text{Ca}^{2+}$ -release channels (125), and  $\text{K}^{+}$  channels (94, 104). It has been reported that the direct effect of  $\text{H}_2\text{O}_2$  on  $\text{K}_{\text{ATP}}$  channels in skeletal muscle is mediated via oxidation of the channel protein (179). It has to be noted that the concentration of  $\text{H}_2\text{O}_2$  used by Weik and Neumcke (179) greatly exceeded those reported in studies where the effect was thought to be indirect (50, 123). Tokube et al. (169) suggested that ROS directly affected the  $\text{K}_{\text{ATP}}$  channel by binding to the ATP-binding site, causing a decrease in the sensitivity of the channel to ATP in the range of 0.2-2 mM, without affecting ADP or glibenclamide binding sites. Indirect effects of ROS on ion transport pathways are mediated via membrane phospholipids. There are several examples where changes in ion transport have been attributed primarily to changes in membrane phospholipids. It has been argued that ROS caused peroxidation of membrane phospholipids and that this led to changes in the  $\text{K}_{\text{ATP}}$  channel (63) and the  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ -ATPase (see Table 7).

The data in Tables 1-9 show that the concentrations of ROS-induced changes are different for ion channels, pumps, and exchangers. It appears that the inhibitory concentrations for ion pumps are less than those required for ion channel inhibition. Thus ion pumps are more sensitive to ROS than ion channels. However, it is not known which of the ion pumps is the primary target. Attempts have been made to determine the primary ion pathway that is affected by ROS from the  $\text{IC}_{50}$  of individual ROS. The data in Tables 7 and 8 show that  $\text{Ca}^{2+}$  uptake is more sensitive to  $\text{H}_2\text{O}_2$  than ouabain-sensitive  $\text{Rb}^{+}$  uptake. However,  $\text{Rb}^{+}$  uptake is more sensitive to  $\text{O}_2^{-}$  than  $\text{Ca}^{2+}$  uptake (42). These findings suggest that the primary ion pump that is affected by ROS depends not only on the type of pump but also on the individual ROS.

## ► MECHANISMS OF ROS-INDUCED MODIFICATIONS

Figure 3 shows the possible molecular targets underlying ROS effects on ion transport mechanisms. These molecular targets include 1) membrane phospholipids, 2) membrane proteins, 3) regulators of ion transport mechanisms, or 4) a combination of these targets.



View larger version (13K):

[\[in this window\]](#)

[\[in a new window\]](#)

Fig. 3. Molecular targets underlying ROS-induced malfunction of transport mechanisms.

### *Oxidation of SH Groups*





ROS that mimic the action of SH-oxidizing agents may act via a different mechanism. It is suggested that oxidation of  $K_{Ca}$  channels by  $H_2O_2$  forms disulfide bonds that differ from those induced by SH oxidation with DTNB and thimerosal (18). There are several examples to support this suggestion. DTDP increased the  $P_o$  of the ATP-sensitive SCl channel (95, 96) and also activated  $H_2O_2$ -induced inhibition of this channel (unpublished observations). Similarly, the inhibitory effects of DTNB on whole cell  $Ca^{2+}$  and  $K^+$  currents in  $\beta$  cells and the effectiveness of  $H_2O_2$  suggest that these known SH-oxidizing agents act differentially (100, 101). It is also possible that oxidizing agents, e.g., oxidized glutathione (GSSH) and DTNB, could have different effects on the same channel (23).

*SH-reducing agents reverse ROS action.* ROS-induced changes that have been reported to be reversed with SH-reducing agents, e.g., DTT, include 1)  $H_2O_2$ -induced increase in  $P_o$  of RyR in both cardiac and skeletal muscle (46, 124), 2)  $H_2O_2$ -induced decline in activity of  $Ca^{2+}$ -activated  $K^+$  channels (18), 3)  $H_2O_2$ -induced depression in the  $Ca^{2+}$  pump (86, 87), 4) UV-C-generated  $\cdot OH$  and peroxy ( $ROO\cdot$ )- or  $H_2O_2$ -induced inhibition of  $Na^+$ - $K^+$ -ATPase (80, 85), and 5)  $H_2O_2$ -induced inhibition of the  $Na^+$ / $Ca^{2+}$  exchanger (88). Cysteine block of ROS-induced inhibition of the SR  $Ca^{2+}$  pump also suggests the involvement of SH groups (164). In addition, ROS-induced mechanical dysfunction, due to impairment of  $Ca^{2+}$ -ATPase, is prevented by SH-reducing DTT (38, 39, 46). It is assumed that SH-modifying agents act on ROS-induced disulfide by dissociating the  $H_2O_2$ -induced disulfide-linked RyR protein complex (45). However, the possibility that DTT has its own effect cannot be ruled out. Cai and Sauvé (18) suggested that oxidation of  $K_{Ca}$  channels by  $H_2O_2$  forms disulfide bonds that differ from those induced by SH oxidation with DTNB and thimerosal.

*Localization of the SH groups for ROS action.* Localization of the SH groups on which ROS action occurs is achieved by using SH-modifying agents that differ in their pharmacological properties. The studies in which the poorly membrane-permeable thimerosal and the charged DTNB oxidizing agents were used suggest that  $H_2O_2$  inhibits  $K_{Ca}$  channels by interacting with SH groups that are localized on the cytoplasmic side of the channel (see Ref. 18). On the other hand, rose bengal, a ROS-generating system that reverses the blocking effect of ryanodine (see Ref. 185), has an action that suggests a competition between ROS and ryanodine on a binding site that contains some SH groups. *N*-(7-dimethylamino-4-methyl-3-coumarinyl)maleimide labeling of cysteine indicates that this binding site (its oxidized SH groups keep the RyR in the active state) is embedded in the membrane away from the cytoplasmic side of the membrane (124). It is not known whether such differences in the localization of SH groups, cytoplasmic vs. internal, may account for differences in the proposed mechanisms of ROS action. The differences in the sensitivity of ion transport pathways to ROS-generating systems may be due to preferential binding of ROS to the SH groups of amino acids in the transport proteins (61). There is also evidence that indicates the presence of different sites underlying ROS-induced modifications in ion transport pathways. The opposite effects of  $H_2O_2$  (inhibition) and DTDP (activation) on the gating of the SCl channel suggest that these oxidizing agents have different binding sites on the channel protein.

Krippeit-Drews et al. (100, 101) reported that DTNB inhibited both  $\text{Ca}^{2+}$  and  $\text{K}_{\text{ATP}}$  currents, whereas  $\text{H}_2\text{O}_2$  had no effect on the  $\text{Ca}^{2+}$  current while it enhanced the  $\text{K}_{\text{ATP}}$  current. These data point to the presence of another mechanism, other than SH oxidization, that may also be responsible for modulating ion channels. The presence of such different mechanisms may explain the opposite effects of  $\text{H}_2\text{O}_2$  (12) and of  $^1\text{O}_2$  and  $\text{O}^-_2$  radicals (69) observed on the RyR  $\text{Ca}^{2+}$ -release channel. There is also evidence that the SH group modulating ATP-sensitive channels may be close to the ATP binding site. ATP inhibition of the  $\text{K}^+$  channel prevents the irreversible inhibitor NEM from reaching critical SH groups (179). Similarly, ATP inhibition of the SCL channel prevents the oxidizing agent DTDP from activating the channel (unpublished observations).

*Other SH-modulated transport proteins.* Some of the ion channels that are modulated by SH reagents have also been modulated by ROS in accordance with the SH hypothesis. One would expect that all ion channels and pumps that are modulated by SH-reducing and SH-oxidizing agents would also be modulated by ROS and the oxidation-reduction state in vivo. However, it should not be assumed that ROS would act in a manner similar to SH-oxidizing agents. As indicated above, there is evidence, contrary to such similarities, pointing to different mechanisms of actions. Some of the ion channels that are modulated by SH reagents, and not yet examined for ROS effects, include fast transient  $\text{K}^+$  ( $I_{\text{K(A)}}$ ) channels (141), diphtheria toxin channels (116), and reduced human skeletal macroscopic  $\text{Cl}^-$  current (hClC-1) (105).

### *Changes in $\text{Ca}^{2+}$ Homeostasis*

Intracellular  $\text{Ca}^{2+}$  is an important second messenger system, and various cells maintain  $\text{Ca}^{2+}$  homeostasis. ROS-induced functional abnormalities in cardiac muscle are thought to be linked to an increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  (see Refs. 49 and 51), which has been confirmed with the fura 2 technique (16, 63). The broad effects of ROS can also be explained in terms of changes in the  $\text{Ca}^{2+}$  second messenger system. In cardiac tissue, the elevation of cytosolic  $\text{Ca}^{2+}$  ( $\text{Ca}^{2+}$  overload) is linked to various functional abnormalities, e.g., contractile dysfunction and ventricular arrhythmia, associated with ROS-induced tissue damage during ischemiareperfusion (51). ROS-induced changes in  $[\text{Ca}^{2+}]_{\text{cyt}}$  homeostasis of muscles in general could be mediated via depression in sarcolemmal  $\text{Ca}^{2+}$ -ATPase, inhibition in SR  $\text{Ca}^{2+}$ -ATPase (Table 7), modification in the gating of SR  $\text{Ca}^{2+}$ -release channels (Table 2), changes in the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (Table 9), or nonspecific  $\text{Ca}^{2+}$  leakage across membranes (see Ref. 161). The changes in  $\text{Ca}^{2+}$  homeostasis need not be directly due to ROS-induced modifications in  $\text{Ca}^{2+}$  pathways but may also arise indirectly via modifications in other ion pathways. Cai and Sauvé (18) have argued that  $\text{H}_2\text{O}_2$  may modulate agonist-induced  $\text{Ca}^{2+}$  influx, activating nitric oxide synthase, which metabolizes L-arginine to citrulline and nitric oxide, indirectly via depolarization in the membrane potential due to  $\text{H}_2\text{O}_2$ -induced inactivation of  $\text{K}_{\text{Ca}}$  channels. The role of HOCl in increasing intracellular  $\text{Ca}^{2+}$  homeostasis (46) is partly due to its effects on both the sarcolemmal  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (88) and the  $\text{Na}^+-\text{K}^+$ -ATPase (103, 114). In fact, some  $\text{Ca}^{2+}$  pathways are ruled out as a cause for changes in  $\text{Ca}^{2+}$  homeostasis. For example, the

irreversible free-radical-induced decrease in  $\text{Ca}^{2+}$  currents in ventricular myocytes suggests that cellular  $\text{Ca}^{2+}$  overload during reperfusion is unlikely to be due to an increase in the sarcolemmal  $\text{Ca}^{2+}$  influx via voltage-gated  $\text{Ca}^{2+}$  channels (48). Regarding the contribution of other  $\text{Ca}^{2+}$  pathways to changes in  $\text{Ca}^{2+}$  homeostasis, Elmoselhi et al. (43) found that the  $\text{Ca}^{2+}$  pump contributing to the  $\text{IP}_3$ -sensitive pool was damaged by  $\text{H}_2\text{O}_2$  and  $\text{O}_2^-$ , whereas the  $\text{Ca}^{2+}$  pump contributing to the  $\text{IP}_3$ -insensitive pool was only damaged by  $\text{H}_2\text{O}_2$ . The  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  channel and a suspected RyR  $\text{Ca}^{2+}$ -release channel are less sensitive than the  $\text{Ca}^{2+}$  pump. Oxidant-induced changes in  $\text{Ca}^{2+}$  homeostasis are also reported to occur in neurons. For example, oxidation enhanced the aggregation of amyloid  $\beta$  protein (36) that forms  $\text{Ca}^{2+}$  channels (3), thus altering  $\text{Ca}^{2+}$  homeostasis to produce neurotoxicity (see Ref. 53).

### *Lipid Peroxidation*

In addition to the direct effects of ROS on ion channels and pumps underlying the transmembrane signaling mechanism (see Ref. 181), ROS alter compartmentation and ionic homeostasis, via membrane phospholipids, leading to alteration in membrane function (16). It is important to distinguish between two possible consequences of ROS-induced lipid peroxidation. The first possibility is that ROS-induced lipid peroxidation leads to a nonspecific leak of some pathway in the lipid itself, which consequently results in a modification of  $\text{Ca}^{2+}$  homeostasis. The second possibility is that ROS-induced lipid peroxidation modifies the physical properties of phospholipids in such a way that some proteins of ion channels, pumps, exchangers, and/or associated proteins that regulate these transport pathways are altered. The first possibility can be ruled out, since there is overwhelming evidence suggesting that ROS induce specific effects on ion transport pathways (see Tables 1-9). The second possibility cannot be ruled out, as it does not exclude the possibility of direct ROS effects on proteins of the ion transport pathways. There is evidence for ROS-induced membrane peroxidation that causes membrane malfunction. Guerra et al. (58) found that anti-lipoperoxidant partially prevented DHF-induced reduction in the DHP binding sites and that the protectant thiourea (an  $\cdot\text{OH}$  scavenger) prevented lipoperoxidative damage (80). It has also been demonstrated that the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is sensitive to lipid composition (10) and is enhanced by increasing cholesterol content (106). Similarly, the  $\text{Na}^+-\text{K}^+-\text{ATPase}$  is activated by fatty acids, acylglycerols, and related amphiphiles (79). It has also been reported that membrane lipid peroxidation by *t*-BHP modified the physiological automaticity by impairing cellular metabolic functions and damaging lipid membrane structure and ion channel proteins (147).

The mechanism of ROS-induced membrane peroxidation involves biochemical changes that alter the physical properties and inactivate membrane-bound enzymes that regulate membrane permeability. Indeed, loss of endothelial cells, which are a major source of reperfusion-generated free radicals, has been found to be associated with increased formation of lipid peroxidation products, such as malondialdehyde and lipid peroxides (see Ref. 98). There is evidence that lipid peroxidation subsequently leads to alterations in  $\text{Ca}^{2+}$  homeostasis (see Ref. 63). For example, *t*-BHP augments and subsequently attenuates  $\text{Ca}^{2+}$  currents in rabbit sinoatrial node and nodal isolated cells. Modification by *t*-BHP of  $\text{Ca}^{2+}$  homeostasis has also been deduced from an increase in resting tension (122). Therefore, lipid peroxidation has been invoked as a mechanism underlying some diseases. For example, Butterfield et al.

(17) reported that  $\beta$ -amyloid peptide free radical fragments initiated synaptosomal lipoperoxidation that has been implicated in Alzheimer's disease.

### *Oxidative Phosphorylation and ATP Levels*

In endothelial cells there is evidence showing that ATP levels decline under conditions of oxidative stress or  $H_2O_2$ -induced inhibition of glucose-dependent pathways of ATP synthesis (68). Obviously, ATP-sensitive, e.g.,  $K_{ATP}$  channels, or ATP-modulated transport pathways, e.g.,  $Ca^{2+}$  and  $Na^+$  pumps, are likely to be modified if the ATP levels are significantly reduced either 1) directly via ROS-induced effects on the metabolism of ATP production or 2) indirectly via ROS-induced splitting of the ATP to ADP and phosphate (see Ref. 169). It has been reported that  $H_2O_2$  inhibits the glycolytic pathway and oxidative phosphorylation (72), causing an increase in the activity of the  $K_{ATP}$  channel (50, 123). However, the relationship between a decrease in ATP levels and cellular dysfunction is not clear (148). There is evidence showing that cytotoxicity is not coupled to ATP levels. For example, desferrioxamine, an iron chelator, and allopurinol and oxypurinol (XO inhibitors) prevent  $H_2O_2$  cytotoxicity but not a decrease in ATP levels in pulmonary endothelial cells (172). Similarly, after ischemia the ATP level recovers on reperfusion, whereas  $Na^+$ - $K^+$ -ATPase and the glycoside binding sites continue to decrease (see Refs. 82 and 90).

At the ion channel level, the X/XO- and  $H_2O_2$ -induced increase in the  $P_o$  of  $K_{ATP}$  channels recorded in the cell-attached configuration results from a reduction in ATP level due to irreversible inhibition of oxidative phosphorylation and glycolysis rather than to a reduction in the channel sensitivity to ATP (51). However, observations similar to those found in the cell-attached configuration (51) and in the inside-out configuration (32) have been attributed to a direct effect on the ATP sensitivity of the channel, thus ruling out inhibition of oxidative phosphorylation and glycolysis (169). These differences are thought to be due to 1) differences in  $Mg^{2+}$  concentration levels, which affect  $K_{ATP}$  channels (169), and 2) differences in ROS-generating systems, i.e., X/XO producing  $O_2^-$  (169) and  $H_2O_2/FeCl_2$  producing  $\cdot OH$  (32).

### *Changes in pH*

It is known that oxidant stress can modify some pH regulatory mechanisms (see Ref. 14). Subsequently, this causes changes in intracellular pH, which can influence various ion transport mechanisms, such as inactivation of enzymes, damage to  $Na^+$ - $K^+$ -ATPase (90), and modification of  $Ca^{2+}$ -release channels (110, 139) and sarcolemmal  $Cl^-$  conductance (142). ROS-induced SR disruption in ischemic myocardium via interaction with  $H^+$  has also been recognized (65). Initially, it was reported that  $O_2^-$  acts as a signal for the increase in intracellular pH (155). Ikebuchi et al. (74) confirmed that HX/XO generating  $O_2^-$  induced an immediate increase in the intracellular pH of human cultured amnion cells. This  $O_2^-$ -induced increase is not mediated via the  $Na^+/H^+$  exchanger as indicated from the ineffectiveness of removing extracellular  $Na^+$  or blocking its pathway with amiloride. Recently, Wu et al. (182) proposed that the

effects of  $\text{H}_2\text{O}_2$  on cultured rat cardiac myoblasts are not mediated through a rise in intracellular  $\text{Ca}^{2+}$  or inhibition of oxidative phosphorylation. They proposed that ROS effects are mediated via induction of intracellular acidification. The mechanism by which ROS induce pH changes is via inhibition of glycolysis and hydrolysis of ATP rather than inhibition of  $\text{Na}^+/\text{H}^+$  and  $\text{Cl}^-/\text{HCO}^-$  exchangers or a  $\text{Na}^+-\text{HCO}_3^-$  cotransporter (182). These authors have shown that, in the cardiac cell model cell line, H9c2, the intracellular production of  $\cdot\text{OH}$  and not  $\text{O}_2^-$  or  $\text{H}_2\text{O}_2$  is the cause of the acidification. Because it is known that acidosis modifies  $\text{Ca}^{2+}$ -release channels (110, 139) and reduces contraction in cardiac and skeletal muscles (44), Wu et al. (182) argued that the small  $\cdot\text{OH}$ -induced acidification (22) in part contributes to the cardiac stunning seen during reperfusion-ischemia by means of either decreasing the sensitivity of the contractile elements to  $\text{Ca}^{2+}$  concentration or reducing  $\text{Ca}^{2+}$  release from the SR. It should be noted that, according to the pH hypothesis, the effects of ROS on ion transport pathways are indirectly mediated via changes in pH. However, single ion channel studies that show the effects of  $\text{H}_2\text{O}_2$  on the RyR  $\text{Ca}^{2+}$ -release channel in artificial bilayer experiments where the pH is constant indicate that  $\text{H}_2\text{O}_2$  directly affects the RyR  $\text{Ca}^{2+}$ -release channel (12).

## ► ROS AS SECOND MESSENGERS IN ION TRANSPORT PATHWAYS

Recent reports suggest that ROS, or at least  $\text{H}_2\text{O}_2$ , may function as second messenger systems. It has been proposed that  $\text{H}_2\text{O}_2$  modulates a complex of heme-linked NADPH oxidase protein coupled to  $\text{K}^+$  channels that function as an oxygen sensor mechanism in airway chemoreceptors of small lung carcinoma cell lines (176). Closure of this  $\text{K}^+$  channel induces membrane depolarization and enhances  $\text{Ca}^{2+}$  influx that could cause the release of transmitters or modification of spike duration and frequency (176). It is assumed that, for  $\text{H}_2\text{O}_2$  to play a second messenger role, a specificity to  $\text{H}_2\text{O}_2$  modulation must be achieved, as well as sufficient concentrations of  $\text{H}_2\text{O}_2$  accumulated, before it is destroyed by  $\text{H}_2\text{O}_2$  scavengers in a highly reduced cellular environment, e.g., the presence of 1 mM GSH (see Refs. 83, 173, 189). Another channel that is modulated by ROS in a second messenger manner is the SCl channel. Pharmacological and biophysical studies indicate the presence of an  $\text{O}_2$ -sensing mechanism (GSH-GSSH) on the SCl channel protein (95) that is also modulated by  $\text{H}_2\text{O}_2$  (unpublished observations). It remains to be seen whether the  $\text{O}_2$ -sensitive  $\text{K}^+$  channel of the arterial chemoreceptor that is modified by low  $\text{PO}_2$  (47) is sensitive to ROS. It has been reported that an anion channel allows  $\cdot\text{O}_2^-$  permeation into human amnion cells, which consequently causes increases in 1) cytosolic pH, 2)  $[\text{Ca}^{2+}]_{\text{cyt}}$ , and 3) release of arachidonate (74). The interaction of ROS with other second messenger systems could also lead to changes in  $\text{Ca}^{2+}$  levels, e.g.,  $\text{H}_2\text{O}_2$ -induced activation of phospholipase  $\text{A}_2$  and arachidonic acid metabolic pathways (21, 22).

## ► CONCLUSIONS

The well-accepted ROS-induced cardiac dysfunction during ischemia and perfusion, cardiomyopathies, neurotoxicity, inflammation, and aging involves the disruption of various ion transport pathways underlying electrophysiological functions. ROS modify ion transport mechanisms either directly via ion transport pathway proteins and/or ion transport regulatory proteins or indirectly via peroxidation of membrane phospholipids. The nature and sequence of events that lead to the disruptions of these ion transport pathways are not fully understood. ROS-induced modification of SH groups on ion transport proteins leads to changes in the homeostasis of  $\text{Ca}^{2+}$ , a major second messenger system, and perhaps other cytosolic factors. The order of potency and the primary mechanism of cell dysfunction for individual ROS are yet to be determined. The potency of individual ROS and the ion transport mechanism that they primarily affect depend on various factors that include types of tissue. It is obvious, therefore, that such understanding is important for the development of specific drugs for individual ion transport proteins. ROS scavengers, e.g., superoxide dismutase and catalase, thiol-disulfide modifying agents, and  $\text{Ca}^{2+}$  channel modulators, are the bases for therapeutic approaches in free radical-induced ischemic and reperfusion myocardial injury. The cloning of ion transport protein isoforms, utilization of specific antibodies and molecular probes, and direct mutations of specific sites, will enable us to characterize the SH-oxidization sites and enhance our understanding of the structure-function relation for individual transport proteins.

▲ [Top](#)  
 ▲ [Abstract](#)  
 ▲ [Introduction](#)  
 • [Conclusions](#)  
 ▼ [References](#)

## ► ACKNOWLEDGEMENTS

I thank Roger McCart for numerous discussions, suggestions, and critical reading of the manuscript. I also acknowledge Drs. Gunasegaaran Karupiah and Mal Rasmussen for commenting on the manuscript.

## ► FOOTNOTES

This research work was supported by National Health and Medical Research Council of Australia Project Grant 970122.

## ► REFERENCES

1. Akbarali, H. I., D. Biecer, S. E. Ohia, and C. R. Triggle. Similarity of relaxations evoked by BRL 34915, pinacidil, and field-stimulation in rat oesophageal tunica muscularis mucosae. *Br. J. Pharmacol.* 95: 519-525, 1988[[Medline](#)].
2. Antolini, M., P. Debetto, L. Trevisi, and S. Luciani. Diamide: positive inotropic effect in isolated atria and inhibition of  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchange in cardiomyocytes. *Pharmacol. Res.* 23: 163-172, 1991[[Medline](#)].
3. Arispe, N., E. Rojas, and H. B. Pollard. Alzheimer's disease amyloid  $\beta$  protein forms calcium channels in bilayer membranes: blockade by tromethamine and aluminum. *Proc. Natl. Acad. Sci. USA* 90:

▲ [Top](#)  
 ▲ [Abstract](#)  
 ▲ [Introduction](#)  
 ▲ [Conclusions](#)  
 • [References](#)

4. **Barrington, P. L.** Interactions of  $H_2O_2$ , EGTA and patch pipette recording methods in feline ventricular myocytes. *J. Mol. Cell. Cardiol.* 26: 557-568, 1994[[Medline](#)].

6. Behl, C., J. B. Davis, R. Lesley, and D. Schubert. Hydrogen peroxide mediates amyloid  $\beta$  protein toxicity. *Cell* 77: 817-827, 1994[Medline].

8. **Beresewicz, A., and M. Horackova.** Alterations in the electrical and contractile behavior of isolated cardiomyocytes by hydrogen peroxide: possible mechanisms. *J. Mol. Cell. Cardiol.* 23: 899-918, 1991 [Medline].

10. **Bersohn, M. M., D. K. Philipson, and R. S. Weiss.** Lysophosphatidylcholine and sodium-calcium exchange in cardiac sarcolemma: comparison with ischemia. *Am. J. Physiol.* 260 (*Cell Physiol.* 29): C433-C438, 1991 [[Medline](#)].

12. **Boraso, A., and J. Williams.** Modification of the gating cardiac sarcoplasmic reticulum  $\text{Ca}^{2+}$ -release channel by  $\text{H}_2\text{O}_2$  and dithiothreitol. *Am. J. Physiol.* 267 (*Heart Circ. Physiol.* 36): H1010-H1016, 1994[[Medline](#)].

14. **Brisseau, G. F., O. Tsai, T. Nordstrom, J. C. Marshall, S. Grinstein, and O. D. Rotstein.** Oxidant stress inhibits pH regulatory mechanisms in murine peritoneal macrophages. *Surgery* 116: 268-274, 1994[[Medline](#)].

16. **Burton, K. P., A. G. Morris, K. D. Massey, L. M. Buja, and H. K. Hagler.** Free radicals alter ionic levels and membrane phospholipids in cultured rat ventricular myocytes. *J. Mol. Cell. Cardiol.* 22: 1035-1047, 1990[[Medline](#)].

17. **Butterfield, D. A., K. Hensley, M. Harris, M. P. Mattson, and J. Carney.**  $\beta$ -Amyloid peptide free radical fragments initiate synaptosomal lipoperoxidation in a sequence-specific fashion: implication to Alzheimer's disease. *Biochem. Biophys. Res. Commun.* 200: 710-715, 1994[[Medline](#)].
18. **Cai, S., and R. Sauvé.** Effects of thiol-modifying agents on a  $K(Ca^{2+})$  channel intermediate conductance in bovine aortic endothelial cells. *J. Membr. Biol.* 158: 137-158, 1997[[Medline](#)].
19. **Carafoli, E., and S. Longoni.** The plasma membrane in the control of signaling function of calcium. In: *Cell Calcium and the Control of Membrane Transport*, edited by I. J. Mandel, and D. C. Eaton. New York: Rockefeller Univ. Press, 1986, p. 21-30.
20. **Cerbai, E., G. Ambrosio, F. Porciatti, M. Chiariello, A. Giotti, and A. Mugelli.** Cellular electrophysiological basis for oxygen radical-induced arrhythmias: a patch-clamp study in guinea-pig ventricular myocytes. *Circulation* 84: 1773-1782, 1991[[Abstract](#)].
21. **Chakraborti, S., S. K. Batabyal, and T. Chakraborti.** Role of hydroxyl radical in stimulation of arachidonic acid release caused by  $H_2O_2$  in pulmonary smooth muscle cells: protective effects of anion channel blocker. *Mol. Cell. Biochem.* 146: 91-98, 1995[[Medline](#)].
22. **Chakraborti, S., and T. Chakraborti.** Down-regulation of protein kinase C attenuates the oxidant hydrogen peroxide-mediated activation of phospholipase  $A_2$  in primary vascular smooth muscle cells. *Cell. Signal.* 7: 75-83, 1995[[Medline](#)].
23. **Chiamvimonvat, N., B. O'Rourke, T. J. Kamp, R. G. Kallen, F. Hofmann, V. Flockerzi, and E. Marban.** Functional consequences of sulfhydryl modification in the pore-forming subunits of cardiovascular  $Ca^{2+}$  and  $Na^{+}$  channels. *Circ. Res.* 76: 325-334, 1995[[Abstract/Full Text](#)].
24. **Coetzee, W. A., H. Ichikawa, and D. J. Hearse.** Oxidant stress inhibits Na-Ca-exchange current in cardiac myocytes: mediation by sulfhydryl groups? *Am. J. Physiol.* 266 (*Heart Circ. Physiol.* 35): H909-H919, 1994[[Medline](#)].
25. **Coetzee, W. A., T. Y. Nakaura, and J.-F. Faivre.** Effects of thiol-modifying agents on  $K_{ATP}$  channels in guinea pig ventricular cells. *Am. J. Physiol.* 269 (*Heart Circ. Physiol.* 38): H1625-H1633, 1995[[Medline](#)].
26. **Coetzee, W. A., and L. H. Opie.** Effects of oxygen free radicals on isolated cardiac myocytes from guinea-pig ventricle: electrophysiological studies. *J. Mol. Cell. Cardiol.* 24: 651-663, 1992[[Medline](#)].
27. **Cohen, M. V.** Free radicals in ischemic and reperfusion myocardial injury: is this the time for clinical trials? *Ann. Intern. Med.* 111: 918-931, 1989[[Medline](#)].
28. **Condrescu, M., G. Chernaya, V. Kalaria, and J. P. Reeves.** Barium influx mediated by the cardiac sodium exchanger in transfected Chinese hamster ovary cells. *J. Gen. Physiol.* 109: 41-51, 1997[[Abstract/Full Text](#)].
29. **Cosentino, F., J. C. Still, and S. Katusic.** Role of superoxide anion in the mediation of endothelium-dependent contraction. *Hypertension* 23: 229-235, 1994[[Abstract](#)].



30. **Davies, K. J. A., A. T. Quintanilha, G. A. Brooks, and L. Parker.** Free radicals and tissue damage produced by exercise. *Biochem. Biophys. Res. Commun.* 107: 1198-1205, 1982[[Medline](#)].
31. **Dawson, V. L., T. M. Dawson, G. R. Uhl, and S. H. Snyder.** Human immunodeficiency virus type 1 coat protein neurotoxicity mediated by nitric oxide in primary cortical cultures. *Proc. Natl. Acad. Sci. USA* 90: 3256-3259, 1993[[Abstract](#)].
32. **Deutsch, N., and J. N. Weiss.** ATP-sensitive  $K^+$  channel modification by metabolic inhibition in isolated guinea-pig ventricular myocytes. *J. Physiol. (Lond.)* 465: 163-179, 1993[[Medline](#)].
33. **DiPolo, R., and L. Beauge.** In squid axons the  $Ca^{2+}_i$  regulatory site of the  $Na^+/Ca^{2+}$  exchange is drastically modified by sulfhydryl blocking agents: evidence that intracellular  $Ca^{2+}_i$  regulatory and transport sites are different. *Biochim. Biophys. Acta* 1145: 75-84, 1993[[Medline](#)].
34. **Dixon, I. M., C. M. Kaneko, T. Hata, V. Panagia, and N. S. Dhalla.** Alterations in cardiac membrane  $Ca^{2+}$  transport during oxidative stress. *Mol. Cell. Biochem.* 99: 125-133, 1990[[Medline](#)].
35. **Downey, A. K.** Free radicals and their involvement during long-term myocardial ischemia and reperfusion. *Annu. Rev. Physiol.* 52: 487-504, 1990[[Medline](#)].
36. **Dryks, T., E. Dyrks, T. Hartmann, C. Masters, and K. Beyreuther.** Amyloidogenicity of  $\beta A4$  and  $\beta A4$ -bearing amyloid protein precursor fragments by metal-catalyzed oxidation. *J. Biol. Chem.* 267: 18210-18217, 1992[[Abstract](#)].
37. **Dupart, F., E. Guillemare, G. Romey, M. Fink, F. Lesage, and M. Lazduski.** Susceptibility of cloned  $K^+$  channels to reactive oxygen species. *Proc. Natl. Acad. Sci. USA* 92: 11796-11800, 1995[[Abstract](#)].
38. **Eley, D. W., J. M. Eley, B. Korecky, and H. Fliss.** Impairment of cardiac contractility and sarcoplasmic reticulum  $Ca^{2+}$  ATPase activity by hypochlorous acid: reversal by dithiothreitol. *Can. J. Physiol. Pharmacol.* 69: 1677-1683, 1991[[Medline](#)].
39. **Eley, D. W., B. Korecky, and H. Fliss.** Dithiothreitol restores contractile function to oxidant-injured cardiac muscle. *Am. J. Physiol.* 257 (Heart Circ. Physiol. 26): H1321-H1325, 1989[[Medline](#)].
40. **Elliott, S. J., D. L. Kunze, and W. P. Schilling.** Oxidant stress alters ion gradients and decreases membrane potential in cultured pulmonary artery endothelial cells (CPAs) (Abstract). *FASEB J.* 5: A656, 1991.
41. **Elliott, S. J., and W. P. Schilling.** Oxidant stress alters  $Na^+$  pump and  $Na^+-K^+-Cl^-$  cotransport activities in vascular endothelial cells. *Am. J. Physiol.* 263 (Heart Circ. Physiol. 32): H96-H102, 1992[[Medline](#)].
42. **Elmoselhi, A. B., A. Butcher, S. E. Samson, and A. K. Grover.** Free radicals uncouple the

- sodium pump in pig coronary artery. *Am. J. Physiol.* 266 (*Cell Physiol.* 35): C720-C728, 1994[[Medline](#)].
43. Elmoselhi, A. B., S. E. Samson, and A. K. Grover. SR  $\text{Ca}^{2+}$  pump heterogeneity: free radicals and  $\text{IP}_3$ -sensitive and -insensitive pools. *Am. J. Physiol.* 271 (*Cell Physiol.* 40): C1652-C1659, 1996[[Medline](#)].
44. Fabiato, A., and F. Fabiato. Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscles. *J. Physiol. (Lond.)* 276: 233-255, 1978[[Medline](#)].
45. Favero, T. G., A. C. Zable, and J. J. Abramson. Hydrogen peroxide stimulates the  $\text{Ca}^{2+}$  release channel from skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 270: 25557-25563, 1995[[Abstract/Full Text](#)].
46. Fukui, K., M. Kaneda, E. Takahashi, M. Washio, and K. Doi. Protective effects of sulfhydryl compounds on HOCl-induced intracellular  $\text{Ca}^{2+}$  increase in single rat ventricular myocytes. *J. Mol. Cell. Cardiol.* 26: 455-461, 1994[[Medline](#)].
47. Ganfornina, M. D., and J. Lopez-Barneo. Gating of  $\text{O}_2$ -sensitive  $\text{K}^+$  channel of arterial chemoreceptor cells and kinetic modifications induced by low  $\text{PO}_2$ . *J. Gen. Physiol.* 100: 427-455, 1992[[Abstract](#)].
48. Gill, J. S., W. J. McKenna, and A. J. Camm. Free radicals irreversibly decrease  $\text{Ca}^{2+}$  currents in isolated guinea-pig ventricular myocytes. *Eur. J. Pharmacol.* 292: 337-340, 1995[[Medline](#)].
49. Goldhaber, J. I. Free radicals enhance  $\text{Na}^+/\text{Ca}^{2+}$  exchange in ventricular myocytes. *Am. J. Physiol.* 271 (*Heart Circ. Physiol.* 40): H823-H833, 1996[[Medline](#)].
50. Goldhaber, J. I., S. Ji, S. T. Lamp, and J. N. Weiss. Effects of exogenous free radicals on electromechanical function and metabolism in isolated rabbit and guinea pig ventricles. Implications for ischemia reperfusion injury. *J. Clin. Invest.* 83: 1800-1809, 1989[[Medline](#)].
51. Goldhaber, J. I., and J. N. Weiss. Oxygen free radicals and cardiac reperfusion abnormalities. *Hypertension* 20: 118-127, 1992[[Abstract](#)].
52. Goldhaber, J. I., and J. N. Weiss. Sodium-calcium exchange is the transient inward current activated by free radicals generating systems in patch-clamped ventricular myocytes (Abstract). *Circulation* 90: I-30, 1994.
53. Goodman, Y., M. R. Steiner, S. M. Steiner, and M. P. Mattson. Nordihydroguaiaretic acid protects hippocampal neurons against amyloid B-peptide toxicity, and attenuates free radical and calcium accumulation. *Brain Res.* 654: 171-176, 1994[[Medline](#)].
54. Greenwald, R. A. Superoxide dismutase and catalase as therapeutic agents for human diseases. A critical review. *Free Radic. Biol. Med.* 8: 201-209, 1990[[Medline](#)].
55. Grover, A. K., and S. E. Samson. Effects of superoxide radical on  $\text{Ca}^{2+}$  pumps of coronary artery.

*Am. J. Physiol.* 255 (*Cell Physiol.* 24): C297-C303, 1988[[Medline](#)].

56. Grover, A. K., S. E. Samson, and V. P. Fomin. Peroxide inactivates calcium pumps in pig coronary artery. *Am. J. Physiol.* 263 (*Heart Circ. Physiol.* 32): H537-H543, 1992[[Medline](#)].

57. Grover, A. K., S. E. Samson, V. P. Fomin, and E. S. Werstiuk. Effects of peroxide and superoxide on coronary artery: ANG II response and sarcoplasmic reticulum  $\text{Ca}^{2+}$  pump. *Am. J. Physiol.* 269 (*Cell Physiol.* 38): C546-C553, 1995[[Medline](#)].

58. Guerra, L., E. Cerbai, S. Gessi, P. A. Borea, and A. Mugelli. The effect of oxygen free radicals on calcium current and dihydropyridine binding sites in guinea-pig ventricular myocytes. *Br. J. Pharmacol.* 118: 1278-1284, 1996[[Medline](#)].

59. Gupta, J. B., and K. Prasad. Mechanism of  $\text{H}_2\text{O}_2$ -induced modulation of airway smooth muscle. *Am. J. Physiol.* 263 (*Lung Cell. Mol. Physiol.* 7): L714-L722, 1992[[Medline](#)].

60. Gusev, G. P., and A. Y. Bogdanova. Evidence for stimulation of K-Cl cotransport system by phenazine methosulfate. *Biochem. Pharmacol.* 43: 2275-2279, 1992[[Medline](#)].

61. Halliwell, B., and M. C. Gutteridge. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem. J.* 219: 1-14, 1984[[Medline](#)].

62. Harvey, W. J., and R. Bolstein. Uncoupling the red cell sodium pump by proteolysis. *J. Biol. Chem.* 261: 1724-1729, 1986[[Abstract](#)].

63. Hayashi, H., H. Miyata, H. Watanabe, A. Kobayashi, and N. Yamazaki. Effects of hydrogen peroxide on action potential and intracellular  $\text{Ca}^{2+}$  concentration of guinea pig heart. *Cardiovasc. Res.* 23: 767-773, 1989[[Medline](#)].

64. Henschke, P. N., and S. J. Elliott. Oxidized glutathione decreases luminal  $\text{Ca}^{2+}$  content of the endothelial cell  $\text{ins}(1,4,5)\text{P}_3$ -sensitive  $\text{Ca}^{2+}$  store. *Biochem. J.* 312: 485-489, 1995[[Medline](#)].

65. Hess, H. L., S. Krause, and H. A. Kontos. Mediation of sarcoplasmic reticulum disruption in the ischaemic myocardium: proposed mechanism by the interaction of hydrogen ions and oxygen free radicals. *Adv. Exp. Med. Biol.* 161: 377-389, 1983[[Medline](#)].

66. Hilgemann, D. W., A. Collins, and S. Matsuoka. Steady-state and dynamic properties of cardiac sodium-calcium exchange. Secondary modulation by cytoplasmic calcium and ATP. *J. Gen. Physiol.* 100: 933-961, 1992[[Abstract](#)].

67. Hilgemann, D. W., S. Matsuoka, G. A. Nagel, and A. Collins. Steady-state and dynamic properties of cardiac sodium-calcium exchange. Sodium-dependent inactivation. *J. Gen. Physiol.* 100: 905-932, 1992[[Abstract](#)].

68. Hinshaw, D. B., J. M. Burger, R. E. Delius, and P. A. Hyslop. Mechanism of protection of oxidant-injured endothelial cells by glutamine. *Surgery* 108: 298-305, 1990[[Medline](#)].

69. **Holmberg, S. R. M., D. Cumming, Y. Kusama, D. J. Hearse, P. A. Poole-Wilson, M. J. Shattock, and A. J. Williams.** Reactive oxygen species modify the structure and function of the cardiac sarcoplasmic reticulum calcium-release channel. *Cardioscience* 2: 19-25, 1991[[Medline](#)].
70. **Huang, W. H., Y. Wang, and A. Askari.** (Na<sup>+</sup> + K<sup>+</sup>)-ATPase: inactivation and degradation induced by oxygen radicals. *Int. J. Biochem.* 24: 621-626, 1992[[Medline](#)].
71. **Huang, W. H., Y. Wang, and A. Askari.** Different sensitivities of the Na<sup>+</sup>/K<sup>+</sup>-ATPase isoforms to oxidants. *Biochim. Biophys. Acta* 1190: 108-114, 1994[[Medline](#)].
72. **Hyslop, P. A., D. B. Hinshaw, W. A. Halsey, I. U. Schraufsaetter, R. D. Sauerheber, R. G. Spragg, J. H. Jackson, and C. G. Cochrane.** Mechanisms of oxidant-mediated cell-injury: the glycolytic and mitochondrial pathways of ADP phosphorylation are major intracellular targets inactivated by hydrogen peroxide. *J. Biol. Chem.* 263: 1665-1675, 1988[[Abstract](#)].
73. **Ichinari, K., M. Kakei, T. Matsuoka, H. Nakashima, and H. Tanaka.** Direct activation of the ATP-sensitive potassium channel by oxygen free radicals in guinea-pig ventricular cells: its potentiation by MgADP. *J. Mol. Cell. Cardiol.* 28: 1867-1877, 1996[[Medline](#)].
74. **Ikebuchi, Y., N. Masumoto, K. Tasaka, K. Koike, K. Kasahara, A. Miyake, and O. Tanizawa.** Superoxide anion increases intracellular pH, intracellular free calcium, and arachidonate release in human amnion cells. *J. Biol. Chem.* 266: 13233-13237, 1991[[Abstract](#)].
75. **Islam, M. S., P. O. Berggren, and P.-O. Larsson.** Sulfhydryl oxidation induces rapid and reversible closure of the ATP-regulated K<sup>+</sup> channel in the pancreatic  $\beta$ -cell. *FEBS Lett.* 319: 128-132, 1993[[Medline](#)].
76. **Islam, M. S., P. Rorsman, and P.-O. Larsson.** Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release in insulin-secreting cells. *FEBS Lett.* 296: 287-291, 1992[[Medline](#)].
77. **Jabr, R. I., and W. C. Cole.** Alteration in electrical activity and membrane currents induced by intracellular oxygen-derived free radical stress in guinea pig ventricular myocytes. *Circ. Res.* 72: 1229-1244, 1993[[Abstract](#)].
78. **Jabr, R. I., and W. C. Cole.** Oxygen-derived radical stress activates nonselective cation current in guinea pig ventricular myocytes. *Circ. Res.* 76: 812-824, 1995[[Abstract/Full Text](#)].
79. **Jackhays, M. G., Z. Xie, Y. Wang, W. H. Wang, and A. Askari.** Activation of Na<sup>+</sup>/K<sup>+</sup>-ATPase by fatty acids, acylglycerols, and related amphiphiles: structure-activity relationship. *Biochim. Biophys. Acta* 1279: 43-48, 1996[[Medline](#)].
80. **Jamme, I., E. Petit, D. Divoux, A. Gerbi, J. M. Maixent, and A. Nouvelot.** Modulation of mouse cerebral Na<sup>+</sup>,K<sup>+</sup>-ATPase activity by oxygen free radicals. *Neuroreport* 7: 333-337, 1995[[Medline](#)].
81. **Janssen, L. J., and J. Wattie.** Non-neurogenic electrically evoked relaxation in canine airway muscle involves action of free radicals on K<sup>+</sup> channels. *J. Pharmacol. Exp. Ther.* 279: 813-821,

1996[Abstract].

82. Jennings, R. B., K. A. Reimer, M. L. Hill, and S. E. Mayer. Total ischemia in dog hearts in vitro. *Circ. Res.* 49: 892-900, 1981[Medline].
83. Ji, L. L., R. Fu, and E. W. Mitchell. Glutathione and antioxidant enzyme in skeletal muscle: effects of fiber type and exercise intensity. *J. Appl. Physiol.* 73: 1854-1859, 1992[Medline].
84. Josephson, R. A., H. S. Silverman, E. G. Lakatta, M. D. Stern, and J. L. Zweier. Study of the mechanisms of hydrogen peroxide and hydroxyl free radical-induced cellular injury and calcium overload in cardiac myocytes. *J. Biol. Chem.* 266: 2354-2361, 1991[Abstract].
85. Kako, J. K., M. Kato, T. Matsuoka, and A. Mustapha. Depression of membrane-bound  $\text{Na}^+\text{-K}^+\text{-ATPase}$  activity induced by free radicals and ischemia of kidney. *Am. J. Physiol.* 254 (Cell Physiol. 23): C330-C337, 1988[Medline].
86. Kaneko, M., R. E. Beamish, and N. S. Dhalla. Depression of heart sarcolemmal  $\text{Ca}^{2+}$  pump activated by oxygen free radicals. *Am. J. Physiol.* 256 (Heart Circ. Physiol. 25): H368-H374, 1989[Medline].
87. Kaneko, M., V. Elimban, and N. S. Dhalla. Mechanism for depression of heart sarcolemmal  $\text{Ca}^{2+}$  pump by oxygen free radicals. *Am. J. Physiol.* 257 (Heart Circ. Physiol. 26): H804-H811, 1989[Medline].
88. Kato, M., and K. J. Kako.  $\text{Na}^+/\text{Ca}^{2+}$  exchange of isolated sarcolemmal membrane: effects of insulin, oxidants and insulin deficiency. *Mol. Cell. Biochem.* 83: 15-25, 1988[Medline].
89. Khananshvil, D. Distinction between the two basic mechanisms of cation transport in the cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchange system. *Biochemistry* 29: 2437-2442, 1990[Medline].
90. Kim, M.-S., and T. Akera.  $\text{O}_2$  free radicals: cause of ischemia-reperfusion injury to cardiac  $\text{Na}^+\text{-K}^+\text{-ATPase}$ . *Am. J. Physiol.* 252 (Heart Circ. Physiol. 21): H252-H257, 1987[Medline].
91. Kloner, R. A., K. Przyklenk, and P. Whittaker. Deleterious effects of oxygen radicals in ischemia/reperfusion. Resolved and unresolved issues. *Circulation* 80: 1115-1127, 1989[Abstract].
92. Koivisto, A., D. Siemen, and J. Nedergaard. Reversible block of the calcium-activated NSC channel in brown fat cells by the sulfhydryl reagents mercury and thimerosal. *Pflügers Arch.* 425: 549-551, 1993[Medline].
93. Koliwad, S. K., S. J. Elliott, and D. L. Kunze. Oxidized glutathione mediates cation channel activation in calf vascular endothelial cells during oxidant stress. *J. Physiol. (Lond.)* 495: 37-49, 1996[Medline].
94. Koong, A. C., A. J. Giaccia, G. M. Hahn, and A. H. Saad. Activation of potassium channels by hypoxia and reoxygenation in the human lung adenocarcinoma cell line A549. *J. Cell. Physiol.* 156:

2613-2619, 1993.

95. **Kourie, J. I.** A redox  $O_2$  sensor modulates the SR  $Ca^{2+}$  counter-current through voltage- and  $Ca^{2+}$ -dependent chloride channels. *Am. J. Physiol.* 272 (*Cell Physiol.* 41): C324-C332, 1997[[Medline](#)].
96. **Kourie, J. I.** ATP-sensitive voltage- and calcium-dependent chloride channels in sarcoplasmic reticulum vesicles from rabbit skeletal muscle. *J. Membr. Biol.* 157: 39-51, 1997[[Medline](#)].
97. **Kourie, J. I.** Chloride channels in the sarcoplasmic reticulum of muscle. *Prog. Biophys. Mol. Biol.* 68: 263-300, 1998.
98. **Kramer, J. H., B. F. Dickens, V. Misik, and W. B. Weglicki.** Phospholipid hydroperoxides are precursors of lipid alkoxyl radicals produced from anoxia/reoxygenated endothelial cells. *J. Mol. Cell. Cardiol.* 27: 371-381, 1995[[Medline](#)].
99. **Kramer, J. H., I. T. Mak, and W. B. Weglicki.** Differential sensitivity of canine cardiac sarcolemmal and microsomal enzymes to inhibition by free radical-induced lipid peroxidation. *Circ. Res.* 55: 120-124, 1984[[Abstract](#)].
100. **Krippeit-Drews, P., S. Britsch, F. Lang, and G. Drews.** Effects of SH-group reagents on  $Ca^{2+}$  and  $K^+$  channel currents of pancreatic  $\beta$ -cells. *Biochem. Biophys. Res. Commun.* 200: 860-866, 1994[[Medline](#)].
101. **Krippeit-Drews, P., F. Lang, D. Häussinger, and G. Drews.**  $H_2O_2$  induced hyperpolarization of pancreatic B-cells. *Pflügers Arch.* 426: 552-554, 1994[[Medline](#)].
102. **Kukreja, R. C., A. A. Kearns, J. L. Zweier, P. Kuppusamy, and M. L. Hess.** Singlet oxygen interaction with  $Ca^{2+}$ -ATPase of cardiac sarcoplasmic reticulum. *Circ. Res.* 69: 1003-1014, 1991[[Abstract](#)].
103. **Kukreja, R. C., A. B. Weaver, and M. L. Hess.** Sarcolemmal  $Na^+$ - $K^+$ -ATPase: inactivation by neutrophil-derived free radicals and oxidants. *Am. J. Physiol.* 259 (*Heart Circ. Physiol.* 28): H1330-H1336, 1990[[Medline](#)].
104. **Kuo, S. S., A. H. Saad, A. C. Koong, G. M. Hahn, and A. J. Giaccia.** Potassium-channel activation in response to low doses of  $\gamma$ -irradiation involves reactive oxygen intermediates in nonexcitatory cells. *Proc. Natl. Acad. Sci. USA* 90: 908-912, 1993[[Abstract](#)].
105. **Kürz, L. L., S. Wagner, A. L. George, Jr., and R. Rüdel.** Probing the major skeletal muscle chloride channel with  $Zn^{2+}$  and other sulfhydryl-reactive compounds. *Pflügers Arch.* 433: 357-363, 1997[[Medline](#)].
106. **Kutryk, M. J. B., and G. N. Pierce.** Stimulation of sodium calcium exchange by cholesterol incorporation into isolated cardiac sarcolemmal vesicles. *J. Biol. Chem.* 263: 13167-13172, 1988[[Abstract](#)].

107. **Lafon-Cazal, M., S. Pietri, M. Culcasi, and J. Bockaert.** NMDA-dependent superoxide production and neurotoxicity. *Nature* 364: 535-537, 1993[[Medline](#)].
108. **Lee, C., and E. Okabe.** Hydroxyl radical-mediated reduction  $\text{Ca}^{2+}$ -ATPase activity of masseter muscle sarcoplasmic reticulum. *Jpn. J. Pharmacol.* 67: 729-734, 1995.
109. **Lytton, J., M. Westlin, and M. R. Hanley.** Thapsigargin inhibits the sarcoplasmic or endoplasmic reticulum Ca-ATPase family of calcium pumps. *J. Biol. Chem.* 266: 17067-17071, 1991[[Abstract](#)].
110. **Ma, J., and J. Zhao.** Highly cooperative hysteric response of skeletal muscle ryanodine receptor to changes in proton concentrations. *Biophys. J.* 67: 626-633, 1994[[Abstract](#)].
111. **MacFarlane, N. G., and D. J. Miller.** Effects of the reactive oxygen species hypochlorous acid and hydrogen peroxide on force production and calcium sensitivity of rat cardiac myofilament. *Pflügers Arch.* 428: 561-568, 1994[[Medline](#)].
112. **Masumoto, N., K. Tasaka, A. Miyake, and O. Tanizawa.** Superoxide anion increases intracellular free calcium in human myometrial cells. *J. Biol. Chem.* 265: 22533-22536, 1990[[Abstract](#)].
113. **Matsuda, H., A. Noma, Y. Kurachi, and H. Irisawa.** Transient depolarization and spontaneous voltage fluctuations in isolated single cells from guinea-pig ventricles. Calcium-mediated membrane potential fluctuations. *Circ. Res.* 51: 142-151, 1982[[Abstract](#)].
114. **Matsuoka, T., T. Yanagishita, and K. J. Kako.** Effects of leukocyte-derived oxidants on sarcolemmal Na,K,ATPase and calcium transport. *Adv. Exp. Med. Biol.* 248: 621-628, 1989[[Medline](#)].
115. **Matsuura, H., and M. J. Shattock.** Membrane potential fluctuations and transient inward currents induced by reactive oxygen intermediates in isolated rabbit ventricular cells. *Circ. Res.* 68: 319-329, 1991[[Abstract](#)].
116. **Mindell, J. A., H. Zhan, P. D. Huynh, R. H. Collier, and A. Finkelstein.** Reaction of diphtheria toxin channels with sulfhydryl-specific reagents: observation of chemical reactions at the single molecule level. *Proc. Natl. Acad. Sci. USA* 91: 5272-5276, 1994[[Abstract](#)].
117. **Mishra, O. P., M. Delivoria-Papadopoulos, G. Cahillane, and C. Waggerle.** Lipid peroxidation as the mechanism of modification of the affinity of the  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase active sites for ATP,  $\text{K}^+$ ,  $\text{Na}^+$  and strophanthidin in vitro. *Neurochem. Res.* 14: 845-851, 1989[[Medline](#)].
118. **Missiaen, L., C. W. Taylor, and M. J. Berridge.** Spontaneous calcium release from inositol triphosphate-sensitive calcium stores. *Nature* 352: 241-244, 1991[[Medline](#)].
119. **Moghadam, H. F., and W. Winlow.** Hydrogen peroxide and nitric oxide decrease  $\text{Ca}^{2+}$  and  $\text{K}^+$  currents in cultured *Lymnaea* neurones (Abstract). *Jpn. J. Physiol.* 42, Suppl. 2: I-7, 1995.
120. **Mount, D. B., R. S. Hoover, and S. C. Hebert.** The molecular physiology of electroneutral cation-chloride cotransport. *J. Membr. Biol.* 158: 177-186, 1997[[Medline](#)].

121. Nakaya, H., Y. Takeda, N. Tohse, and M. Kanno. Mechanism of the membrane depolarization induced by oxidative stress in guinea-pig ventricular cells. *J. Mol. Cell. Cardiol.* 24: 523-534, 1992[[Medline](#)].
122. Nakaya, H., N. Tohse, and M. Kanno. Electrophysiological derangements induced by lipid peroxidation in cardiac tissue. *Am. J. Physiol.* 253 (*Heart Circ. Physiol.* 22): H1089-H1097, 1987[[Medline](#)].
123. Nakazaki, M., M. Kakei, N. Koriyama, and H. Tanaka. Involvement of ATP-sensitive  $K^+$  channels in free radical-mediated inhibition of insulin secretion in rat pancreatic  $\beta$ -cells. *Diabetes* 44: 878-883, 1995[[Medline](#)].
124. Oba, T., M. Koshita, and M. Yamaguchi.  $H_2O_2$  modulates twitch tension and increases  $P_o$  of  $Ca^{2+}$  release channel in frog skeletal muscle. *Am. J. Physiol.* 271 (*Cell Physiol.* 40): C810-C818, 1996[[Medline](#)].
125. Oba, T., M. Yamaguchi, S. Wang, and J. D. Johnson. Modulation of the  $Ca^{2+}$  channel voltage sensor and excitation-contraction coupling by silver. *Biophys. J.* 63: 1416-1420, 1992[[Abstract](#)].
126. Opie, L. H. Reperfusion injury and its pharmacologic modification. *Circulation* 80: 1049-1062, 1989[[Abstract](#)].
127. Pallandi, R. T., M. A. Perry, and T. J. Campbell. Proarrhythmic effects of an oxygen-derived free radical generating system on action potentials recorded from guinea pig ventricular myocardium: a possible cause of reperfusion-induced arrhythmias. *Circ. Res.* 61: 50-54, 1987[[Abstract](#)].
128. Park, M. K., S. H. Lee, S. J. Lee, W. K. Ho, and Y. E. Earm. Different modulation of  $Ca^{2+}$ -activated  $K^+$  channels by the intracellular redox potential in pulmonary and ear arterial smooth muscle cells of the rabbit. *Pflügers Arch.* 430: 308-314, 1995[[Medline](#)].
129. Parker, J. C. Diamide stimulates calcium-sodium exchange in dog red blood cells. *Am. J. Physiol.* 253 (*Cell Physiol.* 22): C580-C587, 1987[[Medline](#)].
130. Phung, C. D., J. A. Ezieme, and J. F. Turrens. Hydrogen peroxide metabolism in skeletal muscle mitochondria. *Arch. Biochem. Biophys.* 315: 479-482, 1994[[Medline](#)].
131. Pierce, G. N., R. Ward, and K. D. Philipson. Role for sulfur-containing groups in  $Na^+$ - $Ca^{2+}$  exchange of cardiac sarcolemmal vesicles. *J. Membr. Biol.* 94: 217-255, 1986[[Medline](#)].
132. Posterino, G. S., and G. D. Lamb. Effects of reducing agents and oxidants on excitation-contraction coupling in skeletal muscle fibres of rat and toad. *J. Physiol. (Lond.)* 496: 809-825, 1996[[Medline](#)].
133. Rack, M., N. Ruby, and C. Waschew. Effects of some chemical reagents on sodium current inactivation in myelinated nerve fibers of the frog. *Biophys. J.* 50: 557-564, 1986[[Abstract](#)].



134. **Ran, S., C. M. Fuller, M. P. Arrate, R. Latorre, and D. J. Benos.** Functional reconstitution of a chloride channel protein from bovine trachea. *J. Biol. Chem.* 267: 20630-20637, 1992[[Abstract](#)].
135. **Reeves, J. P., C. A. Bailey, and C. C. Hale.** Redox modification of sodium-calcium exchange activity in cardiac sarcolemmal vesicles. *J. Biol. Chem.* 26: 4948-4955, 1986.
136. **Reid, M. B., F. A. Khawli, and M. R. Moody.** Reactive oxygen in skeletal muscle. III. Contractility of unfatigued muscle. *J. Appl. Physiol.* 75: 1081-1087, 1993[[Medline](#)].
137. **Rial, E., I. Arechaga, E. Sainz de La Maza, and D. G. Nicholls.** Effect of hydrophobic sulphhydryl reagents on the uncoupling protein and inner-membrane anion channel of brown-adipose-tissue mitochondria. *Eur. J. Biochem.* 182: 187-193, 1989[[Abstract](#)].
138. **Rohn, T. T., T. R. Hinds, and F. F. Vincenzi.** Inhibition of the Ca pump of intact red blood cells by *t*-butyl hydroperoxide: importance of glutathione peroxidase. *Biochim. Biophys. Acta* 1153: 67-76, 1993[[Medline](#)].
139. **Rousseau, E., and J. Pinkos.** pH modulates conducting and gating behaviour of single calcium release channels. *Pflügers Arch.* 415: 645-647, 1990[[Medline](#)].
140. **Rowe, G. T., N. H. Manson, M. Caplan, and M. L. Hess.** Hydrogen peroxide and hydroxyl radical mediation of activated leukocyte depression of cardiac sarcoplasmic reticulum. *Circ. Res.* 53: 584-591, 1983[[Abstract](#)].
141. **Ruppertsberg, J. P., M. Stocker, O. Pongs, S. H. Heinemann, R. Frank, and M. Koenen.** Regulation of fast inactivation of cloned mammalian  $I_{K(A)}$  channels by cysteine oxidation. *Nature* 352: 711-714, 1991[[Medline](#)].
142. **Rychkov, G. Y., M. Pusch, D. St, J. Astill, M. L. Roberts, T. J. Jentsch, and A. B. Bretag.** Concentration and pH dependence of skeletal muscle chloride channel CIC-1. *J. Physiol. (Lond.)* 497: 423-435, 1996[[Medline](#)].
143. **Saito, M., I. Hisatome, S. Nakajima, and R. Sato.** Possible mechanism of oxygen free radical production by human eosinophils mediated by  $K^+$  channel activation. *Eur. J. Pharmacol.* 291: 217-219, 1995[[Medline](#)].
144. **Sakai, H., and N. Takeguchi.** A GTP-binding protein inhibits a gastric housekeeping chloride channel via intracellular production of superoxide. *J. Biol. Chem.* 269: 23426-23430, 1994[[Abstract](#)].
145. **Salama, G., J. J. Abramson, and G. K. Pike.** Sulfhydryl reagents trigger  $Ca^{2+}$  release from the sarcoplasmic reticulum of skinned rabbit psoas fibers. *J. Physiol. (Lond.)* 454: 389-420, 1992[[Medline](#)].
146. **Satoh, N., M. Nishimura, H. Tanaka, N. Homma, and Y. Watanabe.** Augmentation and subsequent attenuation of  $Ca^{2+}$  current due to lipid peroxidation of the membrane caused by *t*-butyl hydroperoxide in rabbit sinoatrial node. *Br. J. Pharmacol.* 98: 712-723, 1989.
147. **Satoh, N., M. Nishimura, H. Tanaka, and Y. Watanabe.** Electrophysiologic alterations in the

- rabbit nodal cells induced by membrane lipid peroxidation. *Eur. J. Pharmacol.* 292: 233-240, 1995[[Medline](#)].
148. Schilling, W. P., and S. J. Elliot.  $\text{Ca}^{2+}$  signaling mechanisms of vascular endothelial cells and their role in oxidant-induced endothelial cell dysfunction. *Am. J. Physiol.* 262 (Heart Circ. Physiol. 31): H1617-H1630, 1992[[Medline](#)].
149. Schlieff, T., and S. Heinemann.  $\text{H}_2\text{O}_2$ -induced chloride currents are indicative of an endogenous  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchange mechanism in *Xenopus* oocytes. *J. Physiol. (Lond.)* 486: 123-130, 1995[[Medline](#)].
150. Schwartz, R. D., P. Skolnick, and S. M. Paul. Regulation of  $\gamma$ -aminobutyric acid/barbiturate receptor-gated chloride ion flux in brain vesicles by phospholipase A2: possible role of oxygen radicals. *J. Neurochem.* 50: 565-571, 1988[[Abstract](#)].
151. Sen, C. K., I. Kolosova, O. Hanninen, and S. N. Orlov. Inward potassium transport systems in skeletal muscle derived cells are highly sensitive to oxidant exposure. *Free Radic. Biol. Med.* 18: 795-800, 1995[[Medline](#)].
152. Shattock, M. J., D. J. Hearse, and H. Matsuura. Ionic currents underlying oxidant stress-induced arrhythmias. In: *Ionic Currents and Ischemia*, edited by J. Vereecke, van, P. P. Bogaert, and F. Verdonck. Louvain, Belgium: Univ. Press, 1990, p. 288-290.
153. Shattock, M. J., and H. Matsuura. Measurement of  $\text{Na}^+$ - $\text{K}^+$  pump current in isolated rabbit ventricular myocytes using the whole-cell voltage-clamp technique. Inhibition of the pump by oxidant stress. *Circ. Res.* 72: 91-101, 1993[[Abstract](#)].
154. Shi, Z. Q., A. J. Davison, and G. F. Tibbits. Effects of active oxygen generated by DTT/ $\text{Fe}^{2+}$  on cardiac  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchange and membrane permeability to  $\text{Ca}^{2+}$ . *J. Mol. Cell. Cardiol.* 21: 1009-1016, 1989[[Medline](#)].
155. Shibamura, M., T. Kuroki, and K. Nose. Superoxide as a signal for increase in intracellular pH. *J. Cell. Physiol.* 136: 379-383, 1988[[Medline](#)].
156. Siczkowski, M., J. E. Davies, and L. L. Ng.  $\text{Na}^+$ - $\text{H}^+$  exchanger isoform 1 phosphorylation in normal wistar-kyoto and spontaneously hypertensive rats. *Circ. Res.* 76: 825-837, 1995[[Abstract/Full Text](#)].
157. Simchowicz, L. Intracellular pH modulates the generation of superoxide radicals by human neutrophils. *J. Clin. Invest.* 76: 1079-1089, 1985[[Medline](#)].
158. Sjodin, B., Y. H. Westing, and F. S. Apple. Biochemical mechanisms for oxygen free radical formation during exercise. *Sports Med.* 10: 236-254, 1990[[Medline](#)].
159. Somlyo, A. P., and A. V. Somlyo. Signal transduction and regulation in smooth muscle. *Nature* 231: 231-236, 1994.
160. Stein, H. J., J. Esplugues, B. J. Whittle, P. Bauerfeind, R. A. Hinder, and A. L. Blum. Direct

- cytotoxic effect of oxygen radicals on the gastric mucosa. *Surgery* 106: 318-323, 1989[[Medline](#)].
161. Stoyanovsky, D. A., G. Salama, and V. E. Kagan. Ascorbate/iron activate  $\text{Ca}^{2+}$ -release channels of skeletal sarcoplasmic reticulum vesicles reconstituted in lipid bilayers. *Arch. Biochem. Biophys.* 308: 214-221, 1994[[Medline](#)].
162. Stuart, J., I. N. Pessah, T. G. Favero, and J. J. Abramson. Photooxidation of skeletal muscle sarcoplasmic reticulum induces rapid calcium release. *Arch. Biochem. Biophys.* 292: 512-521, 1991.
163. Sun, A. Y. The effect of lipooxygenase on synaptosomal  $(\text{Na}^+ \text{K}^+)$ -ATPase isolated from the cerebral cortex of squirrel monkey. *Biochim. Biophys. Acta* 266: 350-360, 1972[[Medline](#)].
164. Suzuki, Y. J., and G. D. Ford. Inhibition of  $\text{Ca}^{2+}$ -ATPase of vascular smooth muscle sarcoplasmic reticulum by reactive oxygen intermediates. *Am. J. Physiol.* 261 (Heart Circ. Physiol. 30): H568-H574, 1991[[Medline](#)].
165. Suzuki, Y. J., and G. D. Ford. Superoxide stimulates  $\text{IP}_3$ -induced  $\text{Ca}^{2+}$  release from vascular smooth muscle sarcoplasmic reticulum. *Am. J. Physiol.* 262 (Heart Circ. Physiol. 31): H114-H116, 1992[[Medline](#)].
166. Tarr, M., E. Arriaga, and D. Valenzano. Progression of cardiac potassium current modification after brief exposure to reactive oxygen. *J. Mol. Cell. Cardiol.* 27: 1099-1109, 1995[[Medline](#)].
167. Tarr, M., and D. Valenzano. Modification of cardiac action potential by photosensitizer-generated reactive oxygen. *J. Mol. Cell. Cardiol.* 21: 539-543, 1989[[Medline](#)].
168. Thastrup, O., P. J. Cullen, B. K. Drobak, M. R. Hanley, and A. P. Dawson. Thapsigargin, a tumor promotor, discharges intracellular  $\text{Ca}^{2+}$  stores by specific inhibition of endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. *Proc. Natl. Acad. Sci. USA* 87: 2466-2470, 1990[[Abstract](#)].
169. Tokube, K., T. Kiyosue, and M. Arita. Opening of cardiac  $\text{K}_{\text{ATP}}$  channel by oxygen free radicals produced by xanthine oxidase reaction. *Am. J. Physiol.* 271 (Heart Circ. Physiol. 40): H478-H489, 1996[[Medline](#)].
170. Trimm, J. I., G. Salama, and J. J. Abramson. Sulfhydryl oxidation induces rapid calcium release from sarcoplasmic reticulum vesicles. *J. Biol. Chem.* 261: 16092-16098, 1986[[Abstract](#)].
171. Ueda, K., S. Shinohara, T. Yagami, K. Asakura, and K. Kawasaki. Amyloid  $\beta$  protein potentiates  $\text{Ca}^{2+}$  influx through L-type voltage-sensitive  $\text{Ca}^{2+}$  channels: a possible involvement of free radicals. *J. Neurochem.* 68: 265-271, 1997[[Abstract](#)].
172. Varani, J., S. H. Phan, D. F. Gibbs, U. S. Ryan, and P. A. Ward.  $\text{H}_2\text{O}_2$ -mediated cytotoxicity of rat pulmonary endothelial cells: changes in adenosine triphosphate and purine products and effects of protective interventions. *Lab. Invest.* 63: 683-689, 1990[[Medline](#)].
173. Vega-Saenz de Miera, E., and B. Rudy. Modulation of  $\text{K}^+$  channels by hydrogen peroxide.

*Biochem. Biophys. Res. Commun.* 196: 1681-1687, 1992.

174. Vinnikova, A. K., R. C. Kukreja, and M. L. Hess. Singlet oxygen-induced inhibition of cardiac sarcolemmal  $\text{Na}^+$ - $\text{K}^+$ -ATPase. *J. Mol. Cell. Cardiol.* 24: 465-470, 1992[[Medline](#)].
175. Vlessis, A. A., P. Muller, D. Bartos, and D. Trunkey. Mechanism of peroxide-induced cellular injury in cultured adult cardiac myocytes. *FASEB J.* 5: 2600-2605, 1990[[Abstract](#)].
176. Wang, D., C. Youngson, V. Wong, H. Yeger, M. C. Dinauer, E. Vega-Saenz de Miera, B. Rudy, and E. Cutz. NADPH-oxidase and a hydrogen peroxide-sensitive  $\text{K}^+$  channel may function as an oxygen sensor complex in airway chemoreceptors and small cell lung carcinoma cell lines. *Proc. Natl. Acad. Sci. USA* 93: 13182-13187, 1996[[Abstract/Full Text](#)].
177. Wang, G. K., M. S. Brodwick, and D. C. Eaton. Removal of sodium channel inactivation in squid axon by the oxidant chloramine-T. *J. Gen. Physiol.* 86: 289-302, 1985[[Abstract](#)].
178. Wang, Z.-W., and M. Kotlikoff. Regulation of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel gating by sulfhydryl redox agents (Abstract). *Biophys. J.* 70: A401, 1996.
179. Weik, R., and B. Neumcke. ATP-sensitive potassium channels in adult mouse skeletal muscle: characterization of the ATP-binding site. *J. Membr. Biol.* 110: 217-226, 1989[[Medline](#)].
180. Weiss, J. H., C. J. Pike, and C. W. Cotman.  $\text{Ca}^{2+}$  channel blockers attenuate  $\beta$ -amyloid peptide toxicity to cortical neurons in culture. *J. Neurochem.* 62: 372-375, 1994[[Abstract](#)].
181. Wesson, D. E., and S. J. Elliott. Xanthine oxidase inhibits transmembrane signal transduction in vascular endothelial cells. *J. Pharmacol. Exp. Ther.* 270: 1197-1207, 1994[[Abstract](#)].
182. Wu, M.-L., K.-L. Tsai, S.-M. Wang, J.-C. Wu, B.-S. Wang, and Y.-T. Lee. Mechanism of hydrogen peroxide and hydroxyl free radical-induced intracellular acidification in cultured rat cardiac myoblasts. *Circ. Res.* 78: 564-572, 1996[[Abstract/Full Text](#)].
183. Xie, Z., H. M. Jack, Y. Wang, S. M. Periyasamy, G. Blanco, W. H. Huang, and A. Askari. Different oxidant sensitivities of the  $\alpha 1$  and  $\alpha 2$  isoforms of  $\text{Na}^+$ / $\text{K}^+$ -ATPase expressed in baculovirus infected insect cells. *Biochem. Biophys. Res. Commun.* 207: 155-159, 1995[[Medline](#)].
184. Xie, Z. J., Y. H. Huang, A. Askari, W. H. Huang, J. E. Klaunig, and A. Askari. Studies on the specificity of the effects of oxygen metabolites on cardiac sodium pump. *J. Mol. Cell. Cardiol.* 22: 911-920, 1990[[Medline](#)].
185. Xiong, H., E. Buck, J. Stuart, I. N. Pessah, G. Salama, and J. J. Abramson. Rose bengal activates the  $\text{Ca}^{2+}$  release channel skeletal muscle sarcoplasmic reticulum. *Arch. Biochem. Biophys.* 292: 522-528, 1992[[Medline](#)].
186. Yasuda, N., Y. Kasuuya, G. Yamada, H. Haama, T. Masaki, and K. Goto. Loss of contractile activity of endothelin-1 induced by electrical field stimulation generated free radicals. *Br. J. Pharmacol.* 113: 21-28, 1994[[Medline](#)].

187. **Zade-Oppen, A. M. M., and P. K. Lauf.** Thiol-dependent passive K:Cl transport in sheep red blood cells. IX. Modulation by pH in the presence and absence of DIDS and the effect of NEM. *J. Membr. Biol.* 118: 143-151, 1990[[Medline](#)].
188. **Zaidi, N. F., C. F. Lagenaur, J. J. Abramson, I. N. Pessah, and G. Salama.** Reactive disulfides trigger  $Ca^{2+}$  release from sarcoplasmic reticulum via an oxidation reaction. *J. Biol. Chem.* 264: 21725-21736, 1989[[Abstract](#)].
189. **Ziegler, D. M.** Role of reversible oxidation-reduction of enzyme thiols-disulfides in metabolic regulation. *Annu. Rev. Biochem.* 54: 305-329, 1985[[Medline](#)].
190. **Zizi, M., M. Forte, E. Blachly-Dyson, and M. Colombini.** NADH regulates the gating of VDAC, the mitochondrial outer membrane channel. *J. Biol. Chem.* 269: 1614-1616, 1994[[Abstract](#)].

Am J Physiol Cell Physiol 275(1):C1-C24  
0002-9513/98 \$5.00

Copyright © 1998 the American Physiological Society

## This article has been cited by other articles:

- Janssen, L. J., Netherton, S. J., Walters, D. K. (2000).  $Ca^{2+}$ -dependent  $K^{+}$  channels and  $Na^{+}$ - $K^{+}$ -ATPase mediate  $H_2O_2$ - and superoxide-induced relaxations in canine trachealis. *J. Appl. Physiol.* 88: 745-752 [[Abstract](#)] [[Full Text](#)]
- Kourie, J. I., Shorthouse, A. A. (2000). Properties of cytotoxic peptide-formed ion channels. *Am. J. Physiol.* 278: 1063C-1087 [[Abstract](#)] [[Full Text](#)]
- Guo, J., Giles, W. R., Ward, C. A. (2000). Effect of hydrogen peroxide on the membrane currents of sinoatrial node cells from rabbit heart. *Am. J. Physiol.* 279: 992H-999 [[Abstract](#)] [[Full Text](#)]
- Weng, T. X., Mo, L., Hellmich, H. L., Yu, A. S. L., Wood, T., Wills, N. K. (2001). Expression and regulation of CIC-5 chloride channels: effects of antisense and oxidants. *Am. J. Physiol.* 280: 1511C-1520 [[Abstract](#)] [[Full Text](#)]

- [Abstract of this Article](#)
- [Reprint \(PDF\) Version of this Article](#)
- Similar articles found in:  
[Am J Physiol Cell Physiol Online](#)  
[PubMed](#)
- [PubMed Citation](#)
- This Article has been cited by:
- Search Medline for articles by:  
[Kourie, J. I.](#)
- Alert me when:  
[new articles cite this article](#)
- [Download to Citation Manager](#)

HOME	TABLE OF CONTENTS	ARTICLE	ARTICLE	ARTICLE	ARTICLE
ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE
ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE
ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE	ARTICLE